

Geometric Packing under Non-uniform Constraints*

Alina Ene[†]

Sariel Har-Peled[‡]

Benjamin Raichel[§]

December 1, 2011

Abstract

We study the problem of discrete geometric packing. Here, given weighted regions (say in the plane) and points (with capacities), one has to pick a maximum weight subset of the regions such that no point is covered more than its capacity. We provide a general framework and an algorithm for approximating the optimal solution for packing in hypergraphs arising out of such geometric settings. Using this framework we get a flotilla of results on this problem (and also on its dual, where one wants to pick a maximum weight subset of the points when the regions have capacities). For example, for the case of fat triangles of similar size, we show an $O(1)$ -approximation and prove that no PTAS is possible.

*The full version of the paper is available from the arXiv [EHR11].

[†]Department of Computer Science; University of Illinois; 201 N. Goodwin Avenue; Urbana, IL, 61801, USA; ene1@uiuc.edu; <http://www.cs.uiuc.edu/~ene1/>. Work on this paper was partially supported by NSF grants CCF-0728782 and CCF-1016684.

[‡]Department of Computer Science; University of Illinois; 201 N. Goodwin Avenue; Urbana, IL, 61801, USA; sariel@uiuc.edu; <http://www.uiuc.edu/~sariel/>. Work on this paper was partially supported by a NSF AF award CCF-0915984.

[§]Department of Computer Science; University of Illinois; 201 N. Goodwin Avenue; Urbana, IL, 61801, USA; raichel2@uiuc.edu; <http://www.cs.uiuc.edu/~raichel2>. Work on this paper was partially supported by a NSF AF award CCF-0915984.

1 Introduction

Motivation and examples. Consider the problem of *obnoxious facility location* [Tam91, Cap99]; that is, you have to place several facilities, but these facilities are undesired (i.e., obnoxious). Facilities of this type include nuclear reactors, wind farms, airports, power plants, factories, prisons, etc. Facilities can also be semi-desirable – a customer might want to have supermarkets close to their home, but they do not want to have too many of them close by as they increase traffic, noise, etc. One natural way to model this geometrically is to associate each obnoxious facility with its region of undesirability. We also have customers (modeled as points), and each customer has a threshold of how many obnoxious facilities it is willing to accept covering it. Different customers may have different thresholds, for example because more affluent people have stronger political power and it is harder to place obnoxious facilities near their homes.

Naturally, if you allow only a single region to cover each customer, then this is a classical packing problem, and much work has been done on packing disks/balls [SMC⁺07]. However, there are many cases where allowing limited interaction between the packed regions is allowed (after all, these facilities are required for modern existence). As a concrete example of this type of problem, consider the placement of radio stations/cellphone towers. While airports allow only very limited levels of interference⁴, higher levels of such interference is acceptable in residential neighborhoods. However, at a certain point there is going to be resistance to placing more wireless towers in residential areas, as these towers are viewed as causing cancer (this fear might be baseless, but it does not change the political reality of the difficulty of placing such towers). On the other hand, there is little resistance to placing such towers along highways in sparsely populated areas.

In this paper, we are interested in the modeling of such problems and in the computation of an efficient approximation to the optimal solution of such problems.

Modeling.

As hinted by the above, perhaps the most natural way to model this problem is as a generalization of the well known independent set problem.


Independent set is a fundamental discrete optimization problem. Unfortunately, it is not only computationally hard, but it is even hard to approximate to within a factor of $n^{1-\varepsilon}$, for any constant ε [Has99] (under the assumption that $\text{NP} \neq \text{P}$). Surprisingly, the problem is considerably easier in some geometric settings. For example, there is a PTAS⁵ [Cha03, EJS05] for the following problem: Given a set of unit disks in the plane, find a maximum cardinality subset of the disks whose interiors are disjoint. Furthermore, a simple local search algorithm yields the desired approximation: For any $\varepsilon > 0$, the local search algorithm that tries to swap subsets of size $O(1/\varepsilon^2)$ yields a $(1 - \varepsilon)$ -approximation in $n^{O(1/\varepsilon^2)}$ time [CH09, CH11].

The discrete independent set problem. In this paper, we consider packing problems in geometric settings that are natural extensions of the geometric independent set problem described above. As a starting point, motivated by practical applications, we consider the discrete version of the geometric independent set problem in which, in addition to a set of weighted regions, we are given a set of points, and the goal is to select a maximum weight subset of the regions so that each point is contained in at most one of the selected regions. We refer to this problem as the *discrete independent set* problem. Chan and Har-Peled [CH11] studied this discrete variant and proved that one can get a good approximation if the union complexity of the regions is small.


⁴See <http://tinyurl.com/7td67v3> for a story of an airport closing down because of radio interference.

⁵Polynomial time approximation scheme.

Note that the discrete independent set problem captures the continuous version of the independent set problem, since we can place a point in each face of the induced arrangement of the given regions. In fact, the discrete version is considerably harder (in some cases) than the continuous variant. The difficulty lies in that several regions forming a valid solution to an instance of a discrete independent set problem may contain a common point that is not part of the set of points given as input; the figure on the right shows an example in which the middle point, marked as a square, is covered twice by the given valid solution.



To illustrate the difference in difficulty, consider the case when the input consists of a set S of segments (in general position) with their endpoints on a circle, such that for every pair of segments, its members intersect. Clearly, in the continuous version, the maximum independent set of segments is a single segment. However, in this case, the discrete version captures the graph independent set problem. More precisely, we can encode any instance of independent set (i.e., a graph $G = (V, E)$) as an instance of this problem as follows. Every vertex $v \in V$ is mapped to a segment s_v of S , and every edge $uv \in E$, is mapped to the point $s_u \cap s_v$ (which is added to a set of points P). Clearly, an independent set of segments of S (in relation to the point set P) corresponds to an independent set in G . That is, the geometric discrete version is sometimes as hard as the graph independent set problem. For example, the figure on the right depicts the resulting instance encoding independent set for $K_{3,3}$.



The packing problem. In this paper, we are interested in the natural extension of the discrete independent set problem to the case where every point has a capacity and might be covered several times (but not exceeding its capacity). The resulting problem has a flavor of a packing problem, and is defined formally as follows.

Problem 1.1 (PACKREGIONS.) *Given a set \mathcal{D} of regions and a set P of points such that each region r has a weight $w(r)$ and each point p has a capacity $\#(p)$, find a maximum weight subset X of the regions such that, for each point p , the number of regions in X that contain p is at most its capacity $\#(p)$.*

We emphasize that different points might have different capacities, which makes the problem considerably more challenging to solve than the unit capacities case (i.e., the discrete independent set problem). We also consider the following dual problem in which the points have weights and the regions have capacities.

Problem 1.2 (PACKPOINTS.) *Given a set \mathcal{D} of regions and a set P of points such that each region r has a capacity $\#(r)$ and each point p has a weight $w(p)$, find a maximum weight subset X of the points such that each region r contains at most $\#(r)$ points of X .*

Hypergraph framework. These two problems can be stated in a unified way in the language of hypergraphs⁶. Given an instance of **PACKREGIONS**, we construct a hypergraph as follows: Each weighted region is a vertex, and all the regions containing a given point of capacity k become a hyperedge (consisting of these regions) of capacity k . A similar reduction works for **PACKPOINTS**, where the given weighted points are the vertices, and each region of capacity k becomes a hyperedge

⁶A *hypergraph* G is a pair (V, E) , where V is a set of vertices and E is a collection of subsets of V which are called *hyperedges*.

of capacity k consisting of all of the points contained in this region. Therefore the previous two problems are special cases of the following problem.

Problem 1.3 (HGRAPHPACKING.) *Given a hypergraph $G = (V, E)$ with a weight function $w(\cdot)$ on the vertices and a capacity function $\#(\cdot)$ on the hyperedges, find a maximum weight subset $X \subseteq V$, such that $\forall f \in E$ we have $|X \cap f| \leq \#(f)$.*

We will be interested primarily in hypergraphs with certain hereditary properties. A hypergraph property is *hereditary* if the sub-hypergraph induced by any subset of the vertices has the property; an example of a hereditary property of hypergraphs is having bounded VC dimension. Roughly, we are interested in hypergraphs having the **bounded growth property**: For any induced sub-hypergraph on t vertices the number of its hyperedges that contain exactly k vertices is near linear in t and its dependency on k is bounded by $2^{O(k)}$, see Definition 2.1. Such hypergraphs arise naturally when considering points and “nice” regions in the plane.

Our results.

- **Main result.** Our main result is an algorithm that provides a good approximation for **HGRAPHPACKING** as a function of the growth of the hypergraph, see Theorem 3.10. Our result can be viewed as an extension of the work of Chan and Har-Peled [CH11] to these considerably more general and intricate settings. For simplicity, we focus on linear weight functions; we show in Appendix A that our main results extends to the case in which the weight function is a non-negative submodular function.
- **Regions with low union complexity.** In Appendix B, we apply our main result to regions that have low union complexity, and we get the following results:
 - (A) If the union complexity of n regions is $O(nu(n))$ then we get an $O(u(n)^{1/\nu})$ -approximation for **PACKREGIONS**, where ν is the minimum capacity of any point in the given instance. (That is, the problem becomes easier as the minimum capacity increases.) For the case where all the capacities are one, this is the discrete independent set problem, and our algorithm specializes to the algorithm of Chan and Har-Peled [CH11], which gives an $O(u(n))$ -approximation.
 - (B) More specifically, we get a constant factor approximation for **PACKREGIONS** if the union complexity of the regions is linear. This holds for (i) fat-triangles of similar size, (ii) unit axis-parallel cubes in 3d, and (iii) pseudo-disks. See Corollary B.3.
 - (C) Similarly, since the union complexity of fat triangles in the plane is $O(n \log^* n)$ [EAS11, AdBES11b], we get an $O((\log^* n)^{1/\nu})$ approximation for such instances of **PACKREGIONS**.
- **Bi-criteria approximation.** Our main result also implies a bi-criteria approximation algorithm. That is, we can improve the quality of the solution, at the cost of potentially violating low capacity regions. Formally, if the input instance $G = (V, E)$ of **HGRAPHPACKING** has at most $F_k(t) = 2^{O(k)}F(t)$ edges of size k when restricted to any subset of t vertices, then for any integer $\phi \geq 1$, our algorithm yields an $(O((F(n)/n)^{1/\phi}), \phi)$ -approximation to the given instance G of **HGRAPHPACKING**. Specifically, the value of the generated solution X is at least $\Omega(\text{opt}/(F(n)/n)^{1/\phi})$, where opt is the value of the optimal solution, and for every hyperedge $f \in E$, we have $|f \cap X| \leq \max(\phi, \#(f))$.
As an example, for any set of n regions in the plane such that the boundaries of any pair of them intersect $O(1)$ times, the above implies that one can get an $(O(n^{1/\phi}), \phi)$ -approximation for **PACKREGIONS**.

- **Axis-parallel boxes.** The union complexity of axis-parallel rectangles can be as high as quadratic, and therefore we cannot immediately apply our main result to get a good approximation. Instead, we decompose the union of axis-parallel rectangles into regions of low union complexity, and this decomposition together with our main result gives us an $O(\log n)$ approximation for instances of **PACKREGIONS** in which the regions are axis-parallel rectangles in the plane (see Lemma B.9). A more involved analysis also applies to the three dimensional case, where we get an $O(\log^3 n)$ approximation for **PACKREGIONS** for axis parallel boxes (see Lemma B.11).
- **Dual problem.** We show in Appendix B.2 that, by standard lifting techniques, we can apply our result for **PACKREGIONS**, where the regions are disks, to the dual problem of **PACKPOINTSINDISKS**. However, for other regions, the dual problem **PACKPOINTS** seems to be more challenging. Specifically, this is true for the case of axis-parallel rectangles. For this case, we first provide a constant factor approximation for *skyline* instances of the problem; a skyline instance is a set of rectangles that lie on the x -axis. Interestingly, if the set of rectangles is defined in relation to a set of points (and each rectangle contains only a few points), then one can define a near-linear (in the number of points) sized set of rectangles such that each original rectangle is the union of two new rectangles. Combining this with the skyline result and a sparsifying technique, we get an $(O(\log n), 2)$ -approximation; that is, every rectangle b contains at most $\max(2, \#(b))$ points of the solution constructed, and the total weight of the solution is $\Omega(\text{opt}/\log n)$ (see Theorem B.21). (Note that, by applying our general framework directly to this setting, we only get an $(O(n^{1/\phi}), \phi)$ -approximation, for any integer $\phi > 0$.)
- **Packing points into fat triangles.** We provide a polylog bi-criteria approximation for the problem of packing points into fat triangles. This requires proving that one can compute, for a given point set, a small number of canonical subsets such that the point set covered by any fat-triangle (if the set is sufficiently small) is the union of a constant number of these canonical subsets. Proving this requires non-trivial modifications of the result of Aronov *et al.* [AES10]. In addition, we show that a measure defined over a fat triangle can be covered by a few fat triangles, each one of them containing only a constant fraction of the original measure. We believe these two results are of independent interest. Plugging this into the machinery previously developed for axis parallel rectangles yields the new approximation algorithm. See Appendix C for details.
- **PTAS for disks and planes.** We adapt the techniques of Mustafa and Ray [MR10] in order to get a PTAS for instances consisting of unweighted disks and unit-capacity points: we lift the problem to 3d, we construct an approximate conflict graph (as done by Mustafa and Ray), and we use a local search algorithm. This result also implies a PTAS for **PACKPOINTS** for unweighted points and uniform capacity halfspaces in \mathbb{R}^3 . See Appendix D for the details.
- **Hardness.** We show some hardness results for our problems. In particular, we show that **PACKPOINTS** for fat triangles in the plane is as hard as independent set in general graphs (see Lemma E.2). We also show that **PACKREGIONS** is APX-hard (and thus there is no PTAS) for similarly sized fat triangles in the plane (thus “matching” the result of Corollary B.3).

Main technical contribution. In addition to the results mentioned above, our work further develops and extends the techniques for rounding linear programming relaxations for geometric packing problems. Our algorithms use the randomized rounding with alteration technique to round a fractional solution rising out of a natural LP relaxation; this technique has been used in much more general settings [Sri01]. The rounding uses the fractional solution to construct a random sample of the regions. The sampled regions might not form a feasible solution and therefore we need to pick a feasible subset of the sample. This step typically involves selecting an ordering in which to consider the sampled regions and greedily picking a feasible subset based on the ordering

[CH09, CCH09, Var10]. The main technical difficulty lies in finding the ordering of the regions. The main idea behind previous approaches is to build a conflict graph and argue that there exists a vertex of low degree; this vertex gives us a region that is a good candidate for the last region in the ordering, and we can recursively consider the remaining regions. We use a similar approach to construct an ordering, but the conflicts that are relevant in our settings are more complicated. For example, in the geometric independent set problem, a conflict is a pair of regions that overlap. However, in our setting, a conflict involves a larger number of regions and therefore we need to consider a conflict hypergraph. We show that there exists a vertex of low degree in this hypergraph and therefore we are able to extend some of the previous approaches to this more general setting. Unsurprisingly, this extension involves an analysis that is considerably more involved.

Previous work. The work that is closest to ours is the paper of Chan and Har-Peled [CH09] which addresses an easier special case of the problems we consider. Fox and Pach [FP11] presented an n^ϵ approximation for independent set for segments in the plane. The usage of LP relaxations for approximating such problems is becoming more popular. In particular, Chalermsook and Chuzhoy [CC09] use a natural LP relaxation to get an $O(\log \log m)$ -approximation for independent set of axis parallel rectangles in the plane. The geometric set cover problem and the more general problem, the geometric set multi-cover problem, have approximation algorithms that use ϵ -nets to round the natural LP relaxation; see [CCH09] and references therein. Chan and Har-Peled [CH09] used local search to get a PTAS for independent set of pseudo-disks. Independently, Mustafa and Ray [MR10] used similar ideas to get a PTAS for hitting set of pseudo-disks in the plane. There is not much work on the hardness of optimization problems in the geometric settings we are interested in. [CC07] shows that the problem of independent set of axis-parallel boxes in three dimensions is APX-hard (the problem is known to be NP-Hard in the plane). See also [GC11, Har09] and references therein for some recent hardness results. Naturally, in non-geometric settings, there is a vast literature on the problems and techniques we use, see [WS11]. As we already mentioned, our algorithms use the randomized rounding with alteration technique to round a fractional solution. This technique was used in [Sri01] to find an approximate solution to packing integer programs (PIPs) of the form $\{\max wx : Ax \leq b, x \in \mathbb{Z}_+^n\}$, where A is a matrix whose entries are either 0 or 1. The approximation guarantee given in [Sri01] is $O(n^{1/B})$, where $B = \min_i b_i$.

Organization. In Section 2 we define the problem and the associated LP relaxation, and describe some basic tools used throughout the paper. In Section 3 we present the approximation algorithm for the hypergraph case. Due to minor space limitations, we move many of the technical parts of the paper to the appendices. In Appendix A, we extend our main result to the case in which the weight function is a submodular function. In Appendix B we present various applications of our main result. In Appendix C we present the algorithm for packing points into fat triangles. In Appendix D we present a PTAS for some restricted cases. In Appendix E we present some hardness results.

2 Preliminaries

For a maximization problem, an algorithm provides an *α -approximation* if it outputs a solution of value at least opt/α , where opt is the value of the optimal solution. An *(α, β) -approximation* algorithm for **HGRAPHPACKING** is an algorithm that returns a (potentially infeasible) solution of value at least opt/α such that each hyperedge f contains at most $\max(\#(f), \beta)$ vertices of the solution.

2.1 LP Relaxation and the Rounding Scheme

We consider the following natural LP relaxation for the **HGRAPHPACKING** problem. For each vertex v , we have a variable x_v with the interpretation that x_v is 1 if v is selected, and 0 otherwise. For each hyperedge f , we have a constraint that enforces that the number of vertices of f that are selected is at most the capacity of f .

$$\begin{aligned} \text{HYPERGRAPH-LP :} \quad \max \quad & \sum_{v \in \mathbf{V}} w_v x_v \\ & \sum_{v \in f} x_v \leq \#(f) \quad \forall f \in \mathbf{E} \\ & 0 \leq x_v \leq 1 \quad \forall v \in \mathbf{V}. \end{aligned}$$

The **energy** of a subset $X \subseteq \mathbf{V}$ is $\mathcal{E}(X) = \sum_{v \in X} x_v$. In the following, \mathcal{E} denotes the **energy** of the LP solution; that is $\mathcal{E} = \mathcal{E}(\mathbf{V}) = \sum_{v \in \mathbf{V}} x_v$. Note that the energy is at most the number of vertices of the hypergraph. Also, we assume that $\mathcal{E} \geq 1$ (which is always true since all the capacities are at least one).

Definition 2.1 Let $\mathbf{G} = (\mathbf{V}, \mathbf{E})$ be a hypergraph. For any integer k , let $F_k(\cdot)$ denote the function $F_k(t) = \max_{X \subseteq \mathbf{V}, |X| \leq t} \left| \left\{ f \mid f \in \mathbf{E} \text{ and } |X \cap f| = k + 1 \right\} \right|$; that is, $F_k(t)$ is the maximum number of hyperedges of size $k + 1$ of a sub-hypergraph of \mathbf{G} that is induced by a subset of at most t vertices. We say that \mathbf{G} has the **bounded growth** property if the following conditions are satisfied:

- (A) There exists a non-decreasing function $\gamma(\cdot)$ such that $F_k(t) \leq 2^{O(k)} t \gamma(t)$ for any k and t .
- (B) There exists a constant c such that $F_k(xt) \leq c F_k(t)$ for any t, k and x such that $1 \leq x \leq 2$.

This notion of bounded growth is a hereditary property of the hypergraph, and it is somewhat similar to the bounds on the size of set systems with bounded VC dimension. Hypergraphs with bounded growth arise naturally in geometric settings.

The minimum capacity of a packing instance is a useful measure of how hard the instance is; formally, the **minimum capacity** of a given instance \mathbf{G} is

$$\nu = \nu(\mathbf{G}) = \min_{f \in \mathbf{E}} \#(f). \tag{1}$$

Let $\mathbf{G} = (\mathbf{V}, \mathbf{E})$ be a hypergraph, and let $X \subseteq \mathbf{V}$ be a subset of its vertices. The sub-hypergraph of \mathbf{G} **induced** by X is $\mathbf{G}_X = \left(X, \left\{ f \cap X \mid f \in \mathbf{E} \right\} \right)$.

For some basic tools about these settings, see Appendix F.1.

3 Approximate packing for hypergraphs

In this section, we present the algorithm for computing a packing for a given hypergraph $\mathbf{G} = (\mathbf{V}, \mathbf{E})$. We assume that $|\mathbf{E}|$ is polynomial in $|\mathbf{V}|$ and that \mathbf{G} has the bounded growth property introduced in Definition 2.1 (properties which both hold for hypergraphs arising out natural geometric settings). Let x be a solution to the HYPERGRAPH-LP relaxation described in Section 2.1.

3.1 The algorithm

We round the fractional solution to an integral solution using a standard randomized rounding with alteration approach. The first step is to choose an appropriate ordering of the vertices. We will see later how to choose a good ordering; for now, we assume that we are given the ordering. The rounding then proceeds in two phases, the **selection phase** and the **alteration phase**. In the selection phase, we pick a random sample \mathcal{C} of the vertices by selecting each vertex v independently

at random with probability x_v/Δ , where Δ is a parameter that we will determine later. In the alteration phase, we pick a subset of \mathcal{C} as follows: We consider the sampled vertices in the order chosen and we add the current vertex to our solution if the resulting solution remains feasible. We say that a vertex is *selected* if it is present in the sample, and we say that it is *accepted* if it is present in the solution. The main insight is that we can take advantage of the bounded growth property of the hypergraph to show that there is an ordering such that each vertex is accepted with constant probability, provided that it is selected. This will immediately imply that the algorithm achieves a $O(\Delta)$ -approximation.

The main challenge is to prove that a good ordering for the alteration phase exists, that is an ordering such that we accept each selected vertex with constant probability. We now proceed to give such a proof. This proof will suggest a natural $O(n^{C+O(1)})$ time brute force algorithm to actually compute this good ordering, where C is the maximum capacity of an edge in the given instance. In Section F.2 we show how one can improve the dependence on C and make the running time polynomial in the input size.

Running Example 3.1 *To keep the presentation accessible, we interpret this algorithm for instances of **PACKREGIONS** in which the regions are disks. Specifically, we are given a weighted set of disks \mathcal{D} and set of points \mathbf{P} with capacities. The hypergraph has a vertex for each disks in \mathcal{D} and a hyperedge for each point $\mathbf{p} \in \mathbf{P}$; the hyperedge $f_{\mathbf{p}}$ consists of the vertices corresponding to all disks of \mathcal{D} that contain \mathbf{p} .*

In this case, the mysterious quantity $F_k(t)$ (see Definition 2.1) is bounded by the number of faces in an arrangement of t disks that have depth exactly $k + 1$. Since the union complexity of t disks is linear, a standard application of the Clarkson technique implies that $F_k(t) = O(kt)$. Thus in this case we have $\gamma(t) = O(1)$.

3.1.1 Constructing a good ordering

Before we describe how to construct a good ordering of the vertices, it is useful to understand what will force a vertex to be rejected in the alteration phase. With this goal in mind, consider an ordering of the vertices. Let \mathcal{C} be a sample of the vertices in \mathbf{V} such that each vertex v is in \mathcal{C} independently at random with probability x_v/Δ . Let v be a vertex in \mathcal{C} . When we consider v in the alteration phase, we will reject v if and only if there exists a hyperedge f of capacity $\#(f)$ such that f contains v and we have already accepted $\#(f)$ vertices of f . The event that we already accepted $\#(f)$ vertices of f is difficult to analyze. However, as we will see, we can settle for a more conservative analysis that upper bounds the probability that v is rejected, given that *all* of the vertices in \mathcal{C} that appear before v in the ordering are accepted. (In the alteration phase, it is possible that not all vertices in \mathcal{C} that appear before v will be accepted, but this can only help us.) Since we are only interested in the event that \mathcal{C} contains $k + 1$ vertices — the vertex v and k other vertices that appear before v in the ordering — that are contained in a hyperedge of capacity k , only the set of vertices that appear before v in the sample matter, and not the actual ordering of the vertices. With this observation in mind, we define a *k -conflict* to be a set of $k + 1$ vertices that are contained in a hyperedge of capacity k . In the following, \mathcal{H}_k denotes the set of all k -conflicts, and $\mathcal{H} = \cup_k \mathcal{H}_k$ denotes the set of all conflicts. We are interested in the probability of the event that all of the vertices of a k -conflict, h , are present in the sample, and we refer to this probability as the *Δ -potential* of the conflict, $\rho_{\Delta}(h)$. For the analysis it will also be useful to define the unscaled version of this quantity that is the probability that all the vertices of a conflict are present given that we sampled each vertex with probability x_v instead of x_v/Δ . We refer to

this quantity as simply the *potential* of the conflict, $\rho(h)$. Formally, we have

$$\rho_{\Delta}(h) = \prod_{v \in h} \frac{x_v}{\Delta} \quad \text{and} \quad \rho(h) = \prod_{v \in h} x_v.$$

Another quantity of interest is the expected number of conflicts in which a vertex v participates, given that v is in the sample. We refer to this quantity as the Δ -*resistance* of a vertex v in a set of vertices $X \subseteq V$, and we use $\eta_{\Delta}(v, X)$ to denote it: $\eta_{\Delta}(v, X) = \frac{\Delta}{x_v} \sum_{h \in \mathcal{H}, h \subseteq X, v \in h} \rho_{\Delta}(h)$.

The ordering. Note that, if the Δ -resistance of v with respect to the set X of vertices that come before it in the ordering is small, the probability of rejecting v is also small. This suggests that the vertex with least resistance (with respect to V) should be the last vertex in the ordering. This gives us the following algorithm for constructing an ordering: We compute the vertex of least resistance and put it last in our ordering (i.e., it is v_n). We then recursively consider the remaining vertices and we compute an ordering for them. In the following, we assume for simplicity that the resulting ordering is v_1, \dots, v_n .

Note that computing the resistance of a vertex by brute force takes $O(n^{C+O(1)})$ time, where C is the maximum capacity of a hyperedge, and therefore this algorithm is not efficient. We give a polynomial time algorithm for constructing the ordering in Section F.2.

3.2 Analysis

Our main insight is that, if the hypergraph satisfies the bounded growth property defined in Definition 2.1, then for any set $X \subseteq V$ there exists a vertex $v \in X$ such that $\eta_{\Delta}(v, X) \leq 1/4$. We prove this below in Section 3.2.2 (see Lemma 3.9). This proof requires that we set $\Delta = \alpha\gamma(\mathcal{E})^{1/\nu}$, where α is some sufficiently large constant. As such, in the remainder of this section we assume $\Delta = \alpha\gamma(\mathcal{E})^{1/\nu}$.

We now show that given $\eta_{\Delta}(v, X) \leq 1/4$, proving the quality of approximation of the algorithm is straightforward.

Lemma 3.2 *Let \mathcal{C} and \mathcal{O} be the set of vertices that were selected and accepted by the algorithm, respectively. For each i , we have $\Pr[v_i \in \mathcal{O} \mid v_i \in \mathcal{C}] \geq 3/4$.*

The proof is in Appendix G.1.

Corollary 3.3 *The total expected weight of the set of vertices output by the algorithm is $\Omega(\text{opt}/\gamma(\mathcal{E})^{1/\nu})$, where opt is the weight of the optimal solution, and ν is the minimum capacity of the given instance. The proof is in Appendix G.2.*

3.2.1 On the expected number of realized conflicts

To analyze the algorithm we need to understand how conflicts might form during its execution, and show that the damage of such conflicts to the generated solution is limited. To this end, consider the quantity

$$F_k(t) = \max_{A \subseteq X, |A| \leq t} \left| \left\{ f \mid f \in \mathbf{E} \text{ and } |A \cap f| = k + 1 \right\} \right|.$$

This is the maximum number of k -conflicts that can be realized for a set of t vertices. The quantity of interest in the following is $\sum_{h \in \mathcal{H}_k} \rho(h)$, as it is the expected number of conflicts that would be realized if we sampled according to the LP solution. Our goal is to prove that this quantity is

bounded by a function of the energy of the LP (the bound will involve the function $F_k(\cdot)$ defined above).

With this goal in mind, we let R be a random sample of X such that each vertex $v \in X$ is in R independently at random with probability $x_v/2$. We stress that R is a random sample that we use for the purposes of defining a quantity \mathcal{M} (i.e., the expected number of conflicts realized in R), and it should not be confused with the random sample \mathcal{C} that is used by the algorithm. In the following, we bound \mathcal{M} from above in Lemma 3.4 and from below in Lemma 3.5. Putting these two bounds together imply the desired bound on $\sum_{h \in \mathcal{H}_k} \rho(h)$.

A conflict $h \in \mathcal{H}$ is **realized** in R if there is a hyperedge $f \in E$ such that $h = f \cap R$ and $|h| = \#(f) + 1$.

The following is similar in spirit to the Clarkson technique (a similar but simpler argument was used by Chan and Har-Peled [CH11]).

Lemma 3.4 *The expected number of k -conflicts realized in R is $\mathcal{M} = O(F_k(\mathcal{E}(X)))$, where R is a random sample of X such that each vertex $v \in X$ is in R independently at random with probability $x_v/2$.*

The proof is in Appendix G.3.

Lemma 3.5 *For each k -conflict h , the probability that h is realized in R is at least $\rho(h)/2(2e)^k$. Therefore the expected number of k -conflicts realized in R is $\mathcal{M} = \Omega\left(\left(\sum_{h \in \mathcal{H}_k, h \subseteq X} \rho(h)\right)/(2e)^k\right)$.*

The proof is in Appendix G.4.

Putting the above two lemmas together, we get the following.

Lemma 3.6 *For any non-negative integer k we have $\sum_{h \in \mathcal{H}_k, h \subseteq X} \rho(h) = O\left((2e)^k F_k(\mathcal{E}(X))\right)$.*

Running Example 3.7 *In our running example, we have that the expected number of k -conflicts that are being realized by a random sample (sampling more or less according to the LP values) is $\sum_{h \in \mathcal{H}_k} \rho(h) = O\left((2e)^k k \mathcal{E}\right)$. This is a hefty quantity, but the key observation is that if we sample according to the LP values scaled down by a large enough constant, then the probability of such a conflict to be realized drops exponentially with k . In particular, for a sufficiently large constant, the expected number of realized k -conflicts in such a sample is going to be $\leq \mathcal{E}/(10 \cdot 2^k)$. Intuitively, this implies that such conflicts can only cause the algorithm to drop very few vertices during the rounding stage, thus guaranteeing a good solution.*

3.2.2 Resistance is futile, if you pick the right vertex

In the following, we consider a subset X of the vertices and we show that there exists a vertex $v \in X$ whose Δ -resistance $\eta_\Delta(v, X)$ is at most $1/4$. Recall that \mathcal{H}_k is the set of all k -conflicts involving vertices in V . We can rewrite the Δ -resistance of v in X as

$$\eta_\Delta(v, X) = \frac{\Delta}{x_v} \sum_{h \in \mathcal{H}_k, h \subseteq X, v \in h} \rho_\Delta(h) = \frac{1}{x_v} \sum_{k=\nu}^{\infty} \frac{1}{\Delta^k} \sum_{h \in \mathcal{H}_k, h \subseteq X, v \in h} \rho(h).$$

As shown in Lemma 3.6, we can relate the total potential of the conflicts of \mathcal{H}_k that are contained in X to the maximum number of k -conflicts contained in a set of at most $\mathcal{E}(X)$ vertices, where $\mathcal{E}(X) = \sum_{v \in X} x_v$.

Recall that the hypergraph has the bounded growth property (see Definition 2.1) and this property is hereditary. Therefore the function $F_k(\cdot)$ in the lemma above has the two properties described in Definition 2.1 and we get the following corollary.

Corollary 3.8 *We have $\sum_{h \in \mathcal{H}_k, h \subseteq X} \rho(h) = O(2^{ck} \mathcal{E}(X) \gamma(\mathcal{E}(X)))$, where c is a constant.*

We can use Corollary 3.8 to complete the proof of Lemma 3.9 as follows.

Lemma 3.9 *Suppose that the hypergraph \mathbf{G} satisfies the bounded growth property (see Definition 2.1). Let $\Delta = \alpha \gamma(\mathcal{E})^{1/\nu}$, where $\alpha > 0$ is a sufficiently large constant and ν is the minimum capacity of the given instance (see Eq. (1)). Then, for any set $X \subseteq \mathbf{V}$, there exists a vertex $v \in X$ such that $\eta_\Delta(v, X) \leq 1/4$.*

Proof: Let $T = \sum_{v \in X} x_v \eta_\Delta(v, X)$. The quantity $T/\mathcal{E}(X)$ is the weighted average of the resistances of the vertices in X , where the weight of a vertex v is $x_v/\mathcal{E}(X)$. Therefore it suffices to show that $T \leq \mathcal{E}(X)/4$, since the minimum resistance is at most the weighted average. We have

$$\begin{aligned} T &= \sum_{k=\nu}^{\infty} \frac{1}{\Delta^k} \sum_{v \in X} \sum_{\substack{h \in \mathcal{H}_k \\ h \subseteq X \\ v \in h}} \rho(h) = \sum_{k=\nu}^{\infty} \frac{k+1}{\Delta^k} \sum_{\substack{h \in \mathcal{H}_k \\ h \subseteq X}} \rho(h) = \sum_{k=\nu}^{\infty} \frac{k+1}{\Delta^k} \sum_{h \in \mathcal{H}_k} \rho(h) \\ &= \sum_{k=\nu}^{\infty} \frac{k+1}{\Delta^k} O(2^{ck} \mathcal{E}(X) \gamma(\mathcal{E}(X))) \leq \mathcal{E}(X) \cdot \underbrace{\beta \sum_{k=\nu}^{\infty} \left(\frac{2^c}{\Delta}\right)^k (k+1) \gamma(\mathcal{E}(X))}_S, \end{aligned}$$

by Corollary 3.8, where β is some constant. Since $\Delta = \alpha \gamma(\mathcal{E})^{1/\nu}$, we have

$$S = \sum_{k=\nu}^{\infty} \beta \left(\frac{2^c}{\alpha}\right)^k (k+1) \left(\frac{1}{\gamma(\mathcal{E})}\right)^{k/\nu} \gamma(\mathcal{E}(X)) \leq \sum_{k=\nu}^{\infty} \beta \left(\frac{2^c}{\alpha}\right)^k (k+1) \left(\frac{1}{\gamma(\mathcal{E})}\right)^{k/\nu} \gamma(\mathcal{E}) \leq \frac{1}{4},$$

In the second to last inequality, we have used the fact that $\gamma(\cdot)$ is non-decreasing. The last inequality follows if we pick α to be a sufficiently large constant. Therefore $T \leq \mathcal{E}(X)/4$, and the lemma follows. \blacksquare

See Appendix F.2 for details of how to improve the running time.

3.3 The result

Theorem 3.10 *Let $\mathbf{G} = (\mathbf{V}, \mathbf{E})$ be a hypergraph with a weight function $w(\cdot)$ on the vertices and a capacity function $\#(\cdot)$ on the edges, such that $|\mathbf{E}|$ is polynomial in $|\mathbf{V}|$ and \mathbf{G} satisfies the bounded growth property (see Definition 2.1). Then we can compute in polynomial time a subset $X \subseteq \mathbf{V}$ of vertices such that no hyperedge f contains more than its capacity $\#(f)$ vertices of X . Furthermore, in expectation, the total weight of the output set is $\Omega(\text{opt}/\gamma(\mathcal{E})^{1/\nu})$, where opt is the weight of the optimal solution, and ν is the minimum capacity of the given instance.*

Consider an integer constant $\phi > 0$, and observe that one can always relax the capacity constraints of a given instance of **HGRAPHPACKING** by replacing all capacities smaller than ϕ by ϕ . Theorem 3.10 thus implies the following.

Corollary 3.11 *Given an instance of **HGRAPHPACKING**, with the bounded growth property, one can compute in polynomial time a $(O(\gamma(\mathcal{E})^{1/\phi}), \phi)$ -approximation to the optimal solution.*

4 Conclusions

In this paper, we presented a general framework for approximating geometric packing problems with non-uniform constraints. We then applied this framework in a systematic fashion to get improved algorithms for specific instances of this problem, many of which required additional non-trivial ideas. There are several special cases of this problem for which we currently do not know any useful approximation; for example, the special case of packing axis-parallel boxes into points, in which the boxes are in four dimensions is still wide open. Making some progress on these special cases is an interesting direction for future work.

Acknowledgments

The authors thank Timothy Chan, Chandra Chekuri, and Esther Ezra for several useful discussions.

References

- [AdBES11a] B. Aronov, M. de Berg, E. Ezra, and M. Sharir. Improved bound for the union complexity of locally fat objects in the plane. manuscript, 2011.
- [AdBES11b] B. Aronov, M. de Berg, E. Ezra, and M. Sharir. Improved bound for the union of fat objects in the plane. manuscript, 2011.
- [AES10] B. Aronov, E. Ezra, and M. Sharir. Small-size ε -nets for axis-parallel rectangles and boxes. *SIAM J. Comput.*, 39(7):3248–3282, 2010.
- [Cap99] P. Cappanera. A survey on obnoxious facility location problems. Technical report, University of Pisa, 1999.
- [CC07] M. Chlebík and J. Chlebíková. The complexity of combinatorial optimization problems on d -dimensional boxes. *SIAM J. Discrete Math.*, 21(1):158–169, 2007.
- [CC09] P. Chalermsook and J. Chuzhoy. Maximum independent set of rectangles. In *Proc. 20th ACM-SIAM Sympos. Discrete Algorithms*, pages 892–901, 2009.
- [CCH09] C. Chekuri, K. L. Clarkson., and S. Har-Peled. On the set multi-cover problem in geometric settings. In *Proc. 25th Annu. ACM Sympos. Comput. Geom.*, pages 341–350, 2009.
- [CH09] T. M. Chan and S. Har-Peled. Approximation algorithms for maximum independent set of pseudo-disks. In *Proc. 25th Annu. ACM Sympos. Comput. Geom.*, pages 333–340, 2009. cs.uiuc.edu/~sariel/papers/08/w_indep.
- [CH11] T. M. Chan and S. Har-Peled. Approximation algorithms for maximum independent set of pseudo-disks. *CoRR*, abs/1103.1431, 2011.
- [Cha03] T. M. Chan. Polynomial-time approximation schemes for packing and piercing fat objects. *J. Algorithms*, 46(2):178–189, 2003.
- [Cla88] K. L. Clarkson. Applications of random sampling in computational geometry, II. In *Proc. 4th Annu. ACM Sympos. Comput. Geom.*, pages 1–11, 1988.
- [CS89] K. L. Clarkson and P. W. Shor. Applications of random sampling in computational geometry, II. *Discrete Comput. Geom.*, 4:387–421, 1989.
- [CVZ11a] C. Chekuri, J. Vondrák, and R. Zenklusen. Submodular function maximization via the multilinear relaxation and contention resolution schemes. In *Proc. 43th Annu. ACM Sympos. Theory Comput.*, 2011. To appear.
- [CVZ11b] C. Chekuri, J. Vondrák, and R. Zenklusen. Submodular function maximization via the multilinear relaxation and contention resolution schemes. *CoRR*, abs/1105.4593, 2011. Preliminary version in STOC 2011.

- [dBCvKO08] M. de Berg, O. Cheong, M. van Kreveld, and M. H. Overmars. *Computational Geometry: Algorithms and Applications*. Springer-Verlag, 3rd edition, 2008.
- [EAS11] E. Ezra, B. Aronov, and M. Sharir. Improved bound for the union of fat triangles. In *Proc. 16th ACM-SIAM Sympos. Discrete Algorithms*, pages 1778–1785, 2011.
- [EHR11] A. Ene, S. Har-Peled, and B. Raichel. Geometric packing under non-uniform constraints. *CoRR*, abs/1107.2949, 2011.
- [EJS05] T. Erlebach, K. Jansen, and E. Seidel. Polynomial-time approximation schemes for geometric intersection graphs. *SIAM J. Comput.*, 34(6):1302–1323, 2005.
- [FP11] J. Fox and J. Pach. Computing the independence number of intersection graphs. In *Proc. 22nd ACM-SIAM Sympos. Discrete Algorithms*, pages 1161–1165, 2011.
- [GC11] E. Grant and T. M. Chan. Exact algorithms and apx-hardness results for geometric set cover. In *Proc. 23rd Canad. Conf. Comput. Geom.*, 2011.
- [Har09] S. Har-Peled. Being fat and friendly is not enough. *CoRR*, abs/0908.2369, 2009.
- [Har11] S. Har-Peled. *Geometric Approximation Algorithms*. Amer. Math. Soc., 2011.
- [Has99] J. Hastad. Clique is hard to approximate within $n^{1-\epsilon}$. *Acta Mathematica*, pages 105–142, 1999.
- [HS11] S. Har-Peled and M. Sharir. Relative (p, ϵ) -approximations in geometry. *Discrete Comput. Geom.*, 45(3):462–496, 2011.
- [Kan91] V. Kann. Maximum bounded 3-dimensional matching is max snp-complete. *Inform. Process. Lett.*, 37:27–35, January 1991.
- [MMP⁺94] J. Matoušek, N. Miller, J. Pach, M. Sharir, S. Sifrony, and E. Welzl. Fat triangles determine linearly many holes. *SIAM J. Comput.*, 23(1):154–169, 1994.
- [MR10] N. H. Mustafa and S. Ray. Improved results on geometric hitting set problems. *Discrete Comput. Geom.*, 44(4):883–895, 2010.
- [SMC⁺07] P. G. Szabó, M. Cs. Markót, T. Csendes, E. Specht, L. G. Casado, and I. Garcãa. *New Approaches to Circle Packing in a Square: With Program Codes (Springer Optimization and Its Applications)*. Springer-Verlag New York, Inc., 2007.
- [Sri01] A. Srinivasan. New approaches to covering and packing problems. In *Proc. 12th ACM-SIAM Sympos. Discrete Algorithms*, pages 567–576, 2001.
- [Tam91] A. Tamir. Obnoxious facility location on graphs. *SIAM J. Discrete Math.*, 4(4):550–567, 1991.
- [Var10] K. Varadarajan. Weighted geometric set cover via quasi-uniform sampling. In *Proc. 42nd Annu. ACM Sympos. Theory Comput.*, pages 641–648, 2010.
- [WS11] D. P. Williamson and D. B. Shmoys. *The Design of Approximation Algorithms*. Cambridge University Press, 2011.

A Contention resolution schemes

Chekuri *et al.* [CVZ11a, CVZ11b] considered a broad class of rounding schemes, which they called *contention resolution schemes* (CR schemes). Informally, the family of all CR schemes consists of all rounding strategies based on randomized rounding with alteration. The precise definition of a CR scheme is the following.

Let N be a finite ground set of size n , and let $f : 2^N \rightarrow \mathbb{R}_+$ be a submodular⁷ set function over N . Let $\mathcal{I} \subseteq 2^N$ be a downward-closed⁸ family of subsets of N . The problem of maximizing $f(X)$ subject to the constraint that $X \in \mathcal{I}$ generalizes the hypergraph packing problem: the function f satisfies $f(X) = \sum_{v \in X} w_v$, and the family \mathcal{I} is the family of all subsets $X \subseteq V$ such that, for each hyperedge f , $|f \cap X| \leq \#(f)$. Let $P_{\mathcal{I}} \subseteq [0, 1]^n$ be a convex relaxation⁹ of the constraints imposed by \mathcal{I} ; the set of all feasible fractional solutions to the HYPERGRAPH-LP relaxation described in Section 2.1 is a convex relaxation for the family of all feasible solutions to the hypergraph packing problem. Let F be the multilinear extension¹⁰ of f . Let x be a feasible solution to the relaxation $\{\max F(x) : x \in P_{\mathcal{I}}\}$. The definition of the multilinear extension F suggests the following natural rounding strategy: given x , we construct a random set $R(x)$ by picking each $i \in N$ independently at random with probability x_i . The expected value of $f(R(x))$ is equal to $F(x)$, but it is unlikely that $R(x)$ is in \mathcal{I} . To address this, we want to remove some elements from $R(x)$ in order to get a subset $I \subseteq R(x)$ such that $I \in \mathcal{I}$. We want this step to have the property that, for each $i \in N$, the probability that i is in I is at least cx_i , for some parameter $c > 0$. Chekuri *et al.* [CVZ11b] call such a rounding strategy a *c-balanced CR scheme for $P_{\mathcal{I}}$* . In certain settings it is convenient to scale the fractional solution; the rounding strategy described above for the hypergraph packing problem is one such example. This motivates the following more general CR scheme.

Definition A.1 ([CVZ11b]) *A (b, c) -balanced CR scheme for $P_{\mathcal{I}}$ is a scheme such that for any $x \in P_{\mathcal{I}}$, the scheme selects an independent subset $I \subseteq R(bx)$ with the following property: $\Pr[i \in I \mid i \in R(bx)] \geq c$ for every element $i \in N$. The scheme is said to be monotone if $\Pr[i \in I \mid R(bx) = R_1] \geq \Pr[i \in I \mid R(bx) = R_2]$ whenever $i \in R_1 \subseteq R_2$. A scheme is said to be strict if $\Pr[i \in I \mid i \in R(bx)] = c$ for every i .*

Chekuri *et al.* showed that, if I is the output of a monotone (b, c) -balanced CR scheme, the expected value of I is at least $c\mathbf{E}[F(bx)]$.

Theorem A.2 ([CVZ11b]) *Let $f : 2^N \rightarrow \mathbb{R}_+$ be a non-negative submodular function and let x be a point in $P_{\mathcal{I}}$, where $P_{\mathcal{I}}$ is a convex relaxation for $\mathcal{I} \subseteq 2^N$. Let $I(x) \in \mathcal{I}$ be the random output of a monotone (b, c) -balanced CR scheme on $x \in P_{\mathcal{I}}$. If f is non-monotone, let us assume in addition that the CR scheme is strict. Then $\mathbf{E}[f(I(x))] \geq c\mathbf{E}[F(bx)]$.*

The rounding scheme described in Section 3.1 is a monotone $(\Delta, 1/4)$ -balanced CR scheme on $x \in P_{\mathcal{I}}$, where $P_{\mathcal{I}}$ is the set of all feasible solutions to the HYPERGRAPH-LP relaxation. Therefore Theorem 3.10 extends to the setting in which the total weight of a set of vertices is a monotone submodular function instead of a linear function. If the weight function is a submodular function that is non-monotone, we cannot use Theorem A.2, since our CR scheme is not strict. Chekuri

⁷A function $f : 2^N \rightarrow \mathbb{R}$ is submodular if $f(A) + f(B) \geq f(A \cap B) + f(A \cup B)$ for any two subsets A, B of N . Additionally, f is monotone if $f(A) \leq f(B)$ for all subsets A, B such that $A \subseteq B$.

⁸A family \mathcal{I} of subsets of N is downward-closed if $B \in \mathcal{I}$ and $A \subseteq B$ then $A \in \mathcal{I}$.

⁹ $P_{\mathcal{I}}$ is the closure of the set of characteristic vectors of the sets in \mathcal{I} under convex combinations.

¹⁰The multilinear extension $F : [0, 1]^{|N|} \rightarrow \mathbb{R}$ of a function $f : 2^N \rightarrow \mathbb{R}$ is the function $F(x) = \sum_{S \subseteq N} f(S) \prod_{i \in S} x_i \prod_{j \notin S} (1 - x_j)$

et al. have shown that, for a monotone CR scheme, one can remove the strictness condition by performing the following pruning operation on the output of the CR scheme. Let I be the output of a (possibly non-strict) (b, c) -balanced CR scheme. We pick a subset $J \subset I$ as follows. We order the elements of the ground set N arbitrarily. We start with $J = I$ and we consider the elements of J in the order that we fixed. When we consider i , we remove i from J if $f(J - \{i\}) \geq f(J)$. Chekuri *et al.* show that this pruning step allows us to remove the strictness condition from Theorem A.2.

Theorem A.3 ([CVZ11b]) *Let $f : 2^N \rightarrow \mathbb{R}_+$ be a non-negative submodular function and let x be a point in $P_{\mathcal{I}}$. Let $I = I(x) \in \mathcal{I}$ be the random output of a monotone (b, c) -balanced CR scheme on $x \in P_{\mathcal{I}}$, and J be a pruned version of I . Then $\mathbf{E}[f(J)] \geq c \mathbf{E}[F(bx)]$.*

Corollary A.4 *Let $G = (V, E)$ be a hypergraph. Let $w : 2^V \rightarrow \mathbb{R}_+$ be a weight function on the vertices that is non-negative and submodular. Let $\#(\cdot)$ be a capacity function on the edges such that $|E|$ is polynomial in $|V|$ and G satisfies the bounded growth property (see Definition 2.1). Then we can compute in polynomial time a subset $X \subseteq V$ of vertices such that no hyperedge f contains more than its capacity $\#(f)$ vertices of X . Furthermore, in expectation, the total weight of the output set is $\Omega(\text{opt}/\gamma(\mathcal{E})^{1/\nu})$, where opt is the weight of the optimal solution, and ν is the minimum capacity of the given instance.*

B Applications

Using our main result (Theorem 3.10), we get several approximation algorithms for the packing problems mentioned in the introduction. We present some of these results here.

B.1 Packing regions with low union complexity

Let \mathcal{D} be a set of n weighted regions in the plane, and let the maximum union complexity of $m \leq n$ objects of \mathcal{D} be $U(m) = mu(m)$. We assume that (i) $U(n)/n = u(n)$ is a non-decreasing function, and (ii) there exists a constant c , such that $U(xr) \leq cU(r)$, for any r and $1 \leq x \leq 2$. We are also given a set of points P , where each point $\mathbf{p} \in P$ is assigned a positive integer $\#(\mathbf{p})$ which is the capacity of \mathbf{p} .

We are interested in solving **PACKREGIONS** (Problem 1.1) for \mathcal{D} and P . Consider the hypergraph G obtained by creating a vertex for each region and a hyperedge for each subset of regions containing a given point of P . Here, $F_k(t)$ is bounded by the number of faces in the arrangement of t regions of depth exactly $k + 1$. The number of such faces can be bounded by the union complexity by a standard application of the Clarkson technique [Cla88, CS89].

Lemma B.1 *Consider a set of regions in the plane such that the boundary of every pair intersects a constant number of times. The number of faces of depth at most $k + 1$ in the arrangement of any subset of these regions of size t is $O(k^2 U(t/k))$.*

Plugging this bound into Theorem 3.10 yields the following result.

Theorem B.2 *Let \mathcal{D} be a set of m weighted regions in the plane such that the union complexity of any t of them is $U(t) = tu(t)$. Let P be a set of n points in the plane, where there is a capacity $\#(\mathbf{p})$ associated with each point $\mathbf{p} \in P$. There is a polynomial time algorithm that computes a subset $\mathcal{O} \subseteq \mathcal{D}$ of regions such that no point $\mathbf{p} \in P$ is contained in more than $\#(\mathbf{p})$ regions of \mathcal{O} . Furthermore, in expectation, the total weight of the output set is $\Omega(\text{opt}/u(\mathcal{E})^{1/\nu})$, where opt is the*

weight of the optimal solution, \mathcal{E} is the energy of the LP solution and ν is the minimum capacity of the given instance.

Alternatively, for any integer constant ϕ , one can get a $(O(u(\mathcal{E})^{1/\phi}), \phi)$ -approximation to the optimal solution for the given instance.

The following results follow from the theorem above.

Corollary B.3 (A) *The union complexity of pseudo-disks and fat triangles of similar size is linear; that is, $U(t) = O(t)$. Therefore we get an $O(1)$ -approximation for **PACKREGIONS** if the regions are fat triangles of similar size, disks, or pseudo-disks.*

(B) *The union complexity of fat triangles is $O(n \log^* n)$ [AdBES11a]. Therefore we get an $O(\log^* n)$ -approximation for **PACKREGIONS** if the regions are (arbitrary) fat triangles.*

(C) *Consider a set of regions in the plane such that any pair of them intersects a constant number of times (e.g., a set of arbitrary triangles). In this case, $U(t) = O(t^2)$ and $u(t) = O(t)$. Therefore, for any integer constant $\phi > 0$, we get an $(O(\mathcal{E}^{1/\phi}), \phi)$ -approximation for instances of **PACKREGIONS** on such regions.*

B.2 Packing halfspaces, rays and disks

Problem B.4 (A) **PACKHALFSPACES**: *Given a weighted set of halfspaces \mathcal{S} and a set of points \mathcal{P} with capacities in \mathbb{R}^3 , find a maximum weight subset \mathcal{O} of \mathcal{S} so that, for each point \mathbf{p} , the number of halfspaces of \mathcal{O} that contains \mathbf{p} is at most $\#(\mathbf{p})$.*

(B) **PACKRAYSINPLANES**: *Given a weighted set of vertical rays \mathcal{R} and a set of planes \mathcal{H} with capacities in \mathbb{R}^3 , find a maximum weight subset \mathcal{O} of \mathcal{R} so that, for each plane \mathbf{h} , the number of rays of \mathcal{O} that intersect \mathbf{h} is at most $\#(\mathbf{h})$.*

(C) **PACKPOINTSINDISKS**: *Given a set \mathcal{D} of disks with capacities and a weighted set \mathcal{P} of points, find a maximum weight subset \mathcal{O} of the points so that each disk $r \in \mathcal{D}$ contains at most $\#(r)$ points of \mathcal{O} .*

Since the union complexity of halfspaces in three dimensions is linear, we get the following from Theorem 3.10 (and the 3d analogue of Lemma B.1).

Corollary B.5 *One can compute, in polynomial time, a constant factor approximation to the optimal solution of the **PACKHALFSPACES** problem.*

Standard point/plane duality implies that the same result holds for the dual problem. Namely, a point (a, b, c) gets mapped to the plane $z = ax + by - c$ and a plane $z = ax + by + c$ gets mapped to the point $(a, b, -c)$. Also, a point lies below a given plane if and only if the dual point of the plane lies below the dual plane of the point. As such, the dual of an instance of **PACKHALFSPACES** is an instance of **PACKRAYSINPLANES** (and vice versa).

Thus, Corollary B.5 implies the following.

Corollary B.6 *One can compute, in polynomial time, a constant factor approximation to the optimal solution of the **PACKRAYSINPLANES** problem.*

Finally, observe that an instance of **PACKPOINTSINDISKS** can be lifted into an instance of **PACKRAYSINPLANES**, by the standard lifting $f(x, y) = (x, y, x^2 + y^2)$, which maps points and disks in the plane to halfspaces and points in three dimensions [dBCvKO08].

Corollary B.7 *One can compute, in polynomial time, a constant factor approximation to the optimal solution to the **PACKPOINTSINDISKS** problem.*

B.3 Axis Parallel Rectangles/Boxes

B.3.1 Packing rectangles (2d)

Problem B.8 (**PACKRECTSINPOINTS.**) *Given a weighted set \mathcal{B} of axis-parallel rectangles in the plane, and a point set \mathcal{P} with capacities, find a maximum weight subset $\mathcal{O} \subseteq \mathcal{B}$, such that, for any $\mathbf{p} \in \mathcal{P}$, the number of rectangles of \mathcal{O} containing \mathbf{p} is at most $\#(\mathbf{p})$.*

Note that the union complexity of a set of rectangles can be quadratic. Hence we cannot simply use Theorem B.2 to get a meaningful approximation. However, by using the standard approach for approximating the independent set of rectangles, one can get a reasonable approximation as the following lemma testifies.

Lemma B.9 *Given an instance $(\mathcal{B}, \mathcal{P})$ of **PACKRECTSINPOINTS** with m rectangles, one can compute, in polynomial time, a subset $\mathcal{O} \subseteq \mathcal{B}$ of total weight $\Omega(\text{opt}/\log m)$ such that no capacity constraint of \mathcal{P} is violated, where opt is the weight of the optimal solution.*

Proof: It is straightforward to verify that a set of rectangles that intersect a common line have linear union complexity. Therefore it follows from Theorem B.2 that we can get a constant factor approximation for such instances. Given an arbitrary set of axis-parallel rectangles, we can reduce it to the case in which all rectangles intersect a common line as follows.

We construct an interval tree on \mathcal{B} . Let $\mathbf{b} \in \mathcal{B}$ be the median rectangle of \mathcal{B} when sorted by left edges. Let ℓ denote the vertical line which passes through the left edge of \mathbf{b} , and let \mathcal{B}_ℓ denote the set of rectangles it intersects. We associate \mathcal{B}_ℓ with the root of our tree, and then recursively build left and right subtrees for the rectangles in $\mathcal{B} \setminus \mathcal{B}_\ell$ that lie to the left or right of ℓ , respectively. The recursion bottoms out once every rectangle has been stabbed by a line. Clearly the depth of the tree is $O(\log m)$, since each time we choose the median line and only recursively continue on those rectangles that do not intersect it. Therefore there exists a level of the range tree that has a solution of weight $\Omega(\text{opt}/\log m)$. The algorithm now considers each level of the tree separately, and for each node at the given level, it constructs an approximate solution for the rectangles associated with the node using the constant factor approximation algorithm guaranteed by Theorem B.2. Next, the algorithm considers the union of all these solutions to form the solution for this level. Since two rectangles associated with two different nodes at the same depth in the range tree do not intersect, the resulting set is a valid solution.

The algorithm returns the best solution found among all the levels. Clearly, its weight is $\Omega(\text{opt}/\log m)$. ■

B.3.2 Packing axis-parallel boxes (3d)

The union complexity of axis-parallel boxes in \mathbb{R}^3 that contain a common point is also linear, and therefore a similar approach as above will enable us to solve the following problem.

Problem B.10 (**PACKBOXESINPOINTS.**) *Given a weighted set \mathcal{B} of axis-parallel boxes in \mathbb{R}^3 , and a point set \mathcal{P} with capacities, find a maximum weight subset $\mathcal{O} \subseteq \mathcal{B}$, such that, for any $\mathbf{p} \in \mathcal{P}$, the number of boxes of \mathcal{O} containing \mathbf{p} is at most $\#(\mathbf{p})$.*

Lemma B.11 *Given an instance $(\mathcal{B}, \mathcal{P})$ of **PACKBOXESINPOINTS** with m boxes, one can compute, in polynomial time, a subset $\mathcal{O} \subseteq \mathcal{B}$ of total weight $\Omega(\text{opt}/\log^3 m)$ such that no capacity constraint of \mathcal{P} is violated, where opt is the weight of the optimal solution.*

Proof: We build a multi-layer interval tree on the boxes. On the top layer, we build a balanced tree on the x -axis projection of the boxes, where a node $v_{x'}$ stores all boxes intersecting the plane $x = x'$. Next, we build for each such node a secondary interval tree on the y -axis projections, and for each node on this secondary data-structure we build a third layer data-structure on the z -axis projections. All the boxes are stored in the nodes of the third layer data-structure.

First, observe that the boxes stored in a node on the third layer, $v_{x'y'z'}$, all contain the point (x', y', z') . Now, the union complexity of axis parallel boxes all sharing a common point is linear. As such, we can apply Theorem 3.10 to compute a packing that does not violate the capacities and is a constant factor approximation to the optimal (on this restricted set of boxes). Now, for each third layer tree, we find the level that contains the best possible combined solution (taking the union of the solutions at a level is valid since there is no point in common to two boxes that are stored in two different nodes at the same level). This assigns values for each node in the secondary tree. Again, we choose for each secondary tree the level with the maximum total weight solution. We assign this value to the corresponding node in the first layer data-structure. Again, we choose the level with the highest possible value. This corresponds to a valid solution that complies with the capacity constraints.

As for the quality of approximation, observe that for a tree at a given layer, at least a logarithmic factor of the remaining weight of the optimal solution is contained in some level of the tree. As such, each time we go down one layer in this data-structure we lose at most a logarithmic factor of the optimal solution, and hence the quality of approximation of this algorithm is $\Omega(\log^3 m)$. ■

Remark B.12 (A) It is natural to ask if this result can be extended to higher dimensions. However, it is easy to see that the union complexity of n axis-parallel boxes in four dimensions that all contain a common point can be quadratic. As such, this approach would fail miserably. We leave the problem of getting a better approximation for this case as an open problem for further research.

(B) The continuous version seems to be considerably easier. For the weighted case (with unit capacities; that is, the independent set variant) an $O(\log^{d-1} m / \log \log m)$ approximation is known [CH09] by solving the two dimensional case, and then using the above interval tree technique to apply it for higher dimensions. For the unweighted continuous case, an $O(\log \log m \log^{d-2} m)$ approximation is known [CC09].


The discrete version is different than the continuous version because, for example, if considering boxes in \mathbb{R}^3 that all intersect the xy -plane, the induced two dimensional instance fails to encode the capacity constraints, as they rises from points in three dimensions that do not lie on this plane (while in the continuous case, it is enough to solve the induced problem in this plane).

B.3.3 Packing points into rectangles

Problem B.13 (**PACKPNTSINRECTS.**) *Given a weighted set \mathcal{P} of points and a set \mathcal{B} of axis-parallel rectangles with capacities in the plane, find a maximum weight subset $\mathcal{O} \subseteq \mathcal{P}$, such that, for any $\mathbf{b} \in \mathcal{B}$, the number of points of \mathcal{O} contained in \mathbf{b} is at most $\#(\mathbf{b})$.*

We first observe that the hypergraph that arises from a instance $\mathcal{G} = (\mathcal{P}, \mathcal{B})$ of **PACKPNTSINRECTS**, might not have the bounded growth property for any reasonable growth function. To see this consider the following example.

Consider two parallel lines in the plane with positive slope. Place $n/2$ points on each line such that all the points on the top line lie above and to the left of all the points on the bottom line. Let the set of rectangles for this instance of **PACKPNTSINRECTS** be all the rectangles which have a point on the top line as their upper left corner and a point on the bottom line as their lower right corner. In this case any subset of $O(t)$ points from the top line and $O(t)$ points from the bottom line induce a set of $O(t^2)$ hyperedges, each of size 2. Therefore, $F_1(t) = \Omega(t^2)$, and hence Theorem 3.10 only gives an $O(\mathcal{E})$ approximation.



Since we cannot hope to apply our main result to the case of **PACKPNTSINRECTS**, we will instead seek a bi-criterion approximation. Our algorithm here is inspired by the work of Ezra *et al.* [AES10] on ε -nets for rectangles. Before tackling this problem, we will first consider an easier variant, which will be useful later in obtaining a bi-criterion approximation. In the following, we call a set of rectangles such that all their (say) bottom edges lies on a common line a **skyline**.

Problem B.14 (**PACKPNTSINSKYLINE**.) *Given a weighted set P of points and a set \mathcal{B} of skyline rectangles with capacities in the plane, find a maximum weight subset $\mathcal{O} \subseteq P$, such that, for any $b \in \mathcal{B}$, the number of points of \mathcal{O} contained in b is at most $\#(b)$.*

Lemma B.15 *Let P be a set of n points in the plane all placed above the x -axis. Let $F_k(n)$ be the maximum number of different subsets of P of size k that are realized by intersecting P with a rectangle whose bottom edge lies on the x -axis. We have that $F_k(n) = O(nk^2)$.*

Proof: Consider a rectangle b with its bottom edge lying on the x -axis, and which contains k points of P . Lower its top edge till it passes through a point of P , and let p denote this point. Similarly, move its left and right edges till they pass through points of P . Let b' be this new canonical rectangle. Now, let i_{left} (resp. i_{right}) be the number of points of P inside b' that are to the left (resp. right) of p . Clearly, $(p, i_{\text{left}}, i_{\text{right}})$ uniquely identifies this canonical rectangle. This implies the claim as $p \in P$, $i_{\text{left}} \leq k$ and $i_{\text{right}} \leq k$, and hence the numbers of such triples is $O(|P|k^2)$. ■

Lemma B.16 *Given an instance of **PACKPNTSINSKYLINE**, one can compute, in polynomial time, an $O(1)$ -approximation to the optimal solution.*

Proof: Consider the associated hypergraph $G = (V, E)$. By Lemma B.15, this hypergraph has the bounded growth property with $F_k(t) = tO(k^2)$ (here $\gamma(t) = 1$). Therefore, the algorithm of Theorem 3.10 provides the required approximation. ■

Lemma B.17 *Given a set P of n points in the plane, and a parameter k , one can compute a set \mathcal{D} of $O(k^2 n \log n)$ axis-parallel rectangles, such that for any axis-parallel rectangle b , if $|b \cap P| \leq k$, then there exists two rectangles $b_1, b_2 \in \mathcal{D}$ such that $(b_1 \cup b_2) \cap P = b \cap P$.*

Furthermore, consider the graph where two points of P are connected if they belong to the same rectangle in \mathcal{D} . Then the number of edges in this graph is $O(nk \log n)$.

Proof: Find a horizontal line ℓ that splits P equally, and compute all the skyline rectangles that contain at most k points of P (that is, compute both the rectangle above and below the line). By Lemma B.15, the number of such rectangles is $O(nk^2)$. Now, recursively compute the rectangle set

for the points above ℓ , and for the points below ℓ . Clearly, the number of rectangles generated is $O(k^2 n \log n)$, and let \mathcal{D} denote the resulting set of rectangles.


Now, consider any axis-parallel rectangle \mathbf{b} such that $|\mathbf{b} \cap \mathbf{P}| \leq k$. If it does not intersect ℓ then by induction it has the desired property. Otherwise, if \mathbf{b} intersects ℓ , then it can be decomposed into two skyline rectangles, each one of them contains at most k points of \mathbf{P} . By construction, for each of these rectangles there is a rectangle in \mathcal{D} that contains exactly the same set of points.

As for the second claim, we apply a similar argument. Consider an edge \mathbf{pq} in this graph that arise because of a top skyline rectangle of ℓ . Furthermore, assume that \mathbf{p} is higher than \mathbf{q} and to its right. Clearly, there are at most k such edges emanating from \mathbf{p} , as the skyline rectangle having \mathbf{p} as its top right corner and having its left edge through \mathbf{q} contains at most k points, and each such rectangle corresponds to a unique edge. As such, we get that the number of edges in the graph is $E(n) = O(nk) + 2T(n/2) = O(nk \log n)$. ■

Remark B.18 A slightly more careful analysis shows that the number of rectangles in the set computed by Lemma B.17 that contain exactly k points is $O(nk \log n)$. This will not be needed for our analysis.

Remark B.19 Consider an instance $\mathbf{G} = (\mathbf{P}, \mathcal{B})$ of **PACKPNTSINRECTS**. Let $\mathbf{G}' = (\mathbf{P}, \mathcal{D})$ be a modified instance of **PACKPNTSINRECTS** where \mathcal{D} is obtained from \mathcal{B} by replacing each rectangle by two new rectangles whose union covers that same set of points. Lemma B.17 guarantees that this can be done such that $|\mathcal{D}| = O(n^3 \log n)$. One might be tempted to believe that we can plug \mathbf{G}' into Theorem 3.10 in order to get a bi-criteria approximation for \mathbf{G} . Unfortunately, this does not work since (as the following example shows) the hypergraph does not have the bounded growth property for any meaningful growth function.

Consider two parallel lines in the plane with positive slope. Place $\Theta(\log n)$ points of \mathbf{P} on each line (or close to the line) such that the points on (or close to) the top line all lie above and to the left of those on the bottom line. The remaining points of \mathbf{P} will all lie below the points on the diagonals. Let T be the interval tree of \mathbf{P} (using vertical split lines). Specifically, the remaining points of \mathbf{P} will be placed such that each point on the top line lies in a different level of T , and all the points on the bottom line lie in the same node as the rightmost point on the top line. More specifically, the leftmost point on the top line will correspond to the root and the points in order from the left to right on the top line will correspond to continually walking down in the tree.



(A \textcircled{u} in the figure represents a cluster of u points close together.) Now let X be the subset of \mathbf{P} which consists of the two set of $\Theta(\log n)$ points on the diagonal lines. Consider the intersection sub-hypergraph induced by X . Suppose that \mathcal{B} has rectangle for every pair of points in \mathbf{P} that can be obtained as the intersection of a rectangle with \mathbf{P} . Then any pair of points from the top and bottom diagonals will correspond to a hyperedge in this induced sub-hypergraph. Therefore, $F_1(\log n) = \Omega(\log^2 n)$, and hence Theorem 3.10 only gives an $O(\mathcal{E})$ approximation.

Since (as the above remark demonstrates) we cannot directly apply Lemma B.17, our approach will be more roundabout. We first show how to solve the independent set variant of our problem

(i.e., unit capacities). Next, we slice the rectangles of the given instance with non-uniform capacities case into subrectangles with unit capacities, and plug it into the above algorithm to get a meaningful approximation.

Lemma B.20 *Given an instance of $G = (V, \mathcal{B})$ of **PACKPNTSINRECTS** with unit capacities, one can compute a subset $X \subseteq V$, such that the total weight of X is $\Omega(\text{opt}/\log \mathcal{E})$ and each rectangle of \mathcal{B} contains at most 2 points of P , where $n = |P|$.*

Proof: We first use Lemma F.2 to sparsify the given instance. We now have a set of $P \subseteq V$ of $t = \Theta(\mathcal{E} \log \mathcal{E})$ points, and an associated fractional solution, such that none of the constraints are violated. The value of the fractional solution on G_P is $\Omega(\text{opt})$, and as such we restrict our search for a solution to P .

Furthermore, we can assume that the value assigned to each point of P by this fractional solution is exactly $1/M$ (we replicate a point i times if it is assigned value i/M), where $M = O(\log \mathcal{E})$. Note, that none of the rectangles of \mathcal{B} contains more than M points of P . In particular, by Lemma B.17, one can build a set of rectangles \mathcal{D} of size $O(M^2 t \log t)$, such that every rectangle of \mathcal{B} can be covered by the union of two rectangles of \mathcal{D} ; formally, for every $b \in \mathcal{B}$ there exists $b_1, b_2 \in \mathcal{D}$ such that $b \cap P = (b_1 \cup b_2) \cap P$. We build a conflict graph G over P connecting two points if (i) they are both contained in a rectangle of \mathcal{B} , and (ii) there is a rectangle of \mathcal{D} that contains them both. By Lemma B.17 this graph has at most $O(Mt \log t) = O(\mathcal{E} \log^3 \mathcal{E})$ edges and $t = \Theta(\mathcal{E} \log \mathcal{E})$ vertices.

We further add edges to G making a clique out of each group of duplicated points that arose from a single given point of P (this is needed since when duplicating the points we perturbed them in order to maintain the implicit general position assumptions of Lemma B.17, and one needs to guarantee that at most one of these copies is picked to the independent set). Now for a point $p \in P$ with LP value x_p , the number of duplicated points is $x_p M$. Hence the number of edges added for these cliques is

$$\sum_{p \in P} \binom{x_p M}{2} \leq \sum_{p \in P} x_p^2 M^2 \leq M^2 \sum_{p \in P} x_p \leq M^2 \mathcal{E} = O(\mathcal{E} \log^2 \mathcal{E}),$$

and hence the number of edges in G overall is $O(\mathcal{E} \log^3 \mathcal{E})$.

It is easy to verify that G has average degree $O(\log^2 \mathcal{E})$, and the total weight of the vertices is $\Theta(\text{opt} \log \mathcal{E})$, as such, by Turán's theorem, one can compute an independent set of vertices in this graph of weight $\Omega(w(P)/(\text{average degree} + 1)) = \Omega(\text{opt}/\log \mathcal{E})$.

Now, it is easy to verify that any rectangle in \mathcal{B} contains at most two points of this independent set. ■

Theorem B.21 *Given an instance of (V, \mathcal{B}) of **PACKPNTSINRECTS** (with arbitrary capacities), one can compute in polynomial time a subset $X \subseteq V$ that is an $(O(\log \mathcal{E}), 2)$ -approximation to the optimal solution.*

Proof: Compute a fractional solution to the given instance. Split each rectangle b with capacity $\#(b)$ into $\lceil \#(b)/3 \rceil$ rectangles, each one containing at most value 4 from the fractional solution (this can be done by sweeping the rectangle from left to right, and splitting it whenever the fractional solution inside the current portion exceeds 3). Consider now a unit capacity instance on the same point set but with these new rectangles. We use Lemma B.20 in order to get an $(O(\log \mathcal{E}), 2)$ -approximation for this new instance.

We now show that this solution we obtained for the unit capacity instance is also an $(O(\log \mathcal{E}), 2)$ -approximation to the original instance. First observe that the LP value on this new instance is

$\Omega(\text{opt})$ (where opt is the LP value of the original instance) since scaling down the fractional solution to original instance by a factor of 4 would be a valid solution to the LP for the new instance (since these newly created rectangles each contained at most 4 from the fractional solution), and hence the weight of the approximation is $\Omega(\text{opt}/\log \mathcal{E})$. Furthermore, we know every rectangle $\mathbf{b} \in \mathcal{B}$ contains at most $2 \lceil \#(\mathbf{b})/3 \rceil \leq \max(2, \#(\mathbf{b}))$ points from this solution, since in the new instance each rectangle from \mathcal{B} was replaced with $\lceil \#(\mathbf{b})/3 \rceil$ rectangles each of which contains at most two points from the computed solution. (Note that the inequality holds since $\#(\mathbf{b})$ has integral value.) ■

C Packing points into fat triangles

In this section, we give a bi-criterion approximation for packing points into a set of α -fat triangles. More precisely, we consider the following problem.

Problem C.1 (**PACKPOINTSINFATTRIANGS.**) *Given a weighted set \mathcal{P} of points and a set \mathcal{T} of α -fat triangles in the plane such that each triangle Δ has a capacity $\#(\Delta)$, find a maximum weight subset $\mathcal{O} \subseteq \mathcal{P}$, such that, for each $\Delta \in \mathcal{T}$, the number of points of \mathcal{O} contained in Δ is at most $\#(\Delta)$.*

The approximation algorithm uses the following building blocks:

- (A) We prove that, for a given point set, there exists a small number of canonical sets such that for any fat triangle that covers at most k points, there exists a constant number of these canonical sets whose union covers exactly the same points. Showing this result is quite technical and requires non-trivial modifications of the work of Aronov *et al.* [AES10] (in particular, their work does not imply this result). This is delegated to Section C.4, see Theorem C.6 for the exact result.
- (B) An algorithm for approximating the unit capacity case. This follows by an algorithm similar to the one in Lemma B.20, see Lemma C.2 for details. Note that this uses the result from (A) to get the required approximation.
- (C) A partition scheme that shows that a fat triangle (with a measure defined over it) can be “partitioned” into $O(k)$ triangles such that any triangle in this partition has measure at most $1/k$; see Lemma C.3.

Putting these components together yields the approximation algorithm; see Theorem C.5 for details.

C.1 The unit capacity case

Lemma C.2 *Given an instance $\mathcal{G} = (\mathcal{V}, \mathcal{T})$ of **PACKPOINTSINFATTRIANGS** with unit capacities, one can compute a subset $X \subseteq \mathcal{V}$ such that the total weight of X is $\Omega(\text{opt}/\log^6 \mathcal{E})$ and each triangle of \mathcal{T} contains at most 9 points of \mathcal{P} .*

Proof: We follow the proof of Lemma B.20. We first use Lemma F.2 to sparsify the given instance. We now have a set of $\mathcal{P} \subseteq \mathcal{V}$ of $t = \Theta(\mathcal{E} \log \mathcal{E})$ points and a corresponding fractional solution that is feasible. The value of the fractional solution on $\mathcal{G}_{\mathcal{P}}$ is $\Omega(\text{opt})$, and as such we restrict our search for a solution to \mathcal{P} .

Furthermore, we can assume that the value assigned to each point of \mathcal{P} by this fractional solution is exactly $1/M$ — we replicate a point i times if it is assigned value i/M — where $M = O(\log \mathcal{E})$. Note that none of the triangles of \mathcal{T} contains more than M points of \mathcal{P} . In particular, by Theorem C.6, one can construct a set \mathcal{Z} of regions of size $O(M^3 t \log^2 t)$ such that, for every

triangle of $\Delta \in \mathcal{T}$, there exists a subset $\{z_1, \dots, z_k\} \subseteq \mathcal{Z}$ of at most 9 regions (i.e. $k \leq 9$) such that $\mathbf{P} \cap \Delta = \mathbf{P} \cap (\cup_{i=1}^k z_i)$. We build a conflict graph G over \mathbf{P} connecting two points if (i) they are both contained in a triangle of \mathcal{T} , and (ii) there is a set of \mathcal{Z} that contains both of them. Since the number of sets in \mathcal{Z} is $O(M^3 t \log^2 t)$, and each such set has size at most M , it follows that the number of edges in the resulting graph G is $O(M^5 t \log^2 t)$, and the number of vertices is $t = \Theta(\mathcal{E} \log \mathcal{E})$ vertices. As in the proof of Lemma B.20, we also add edges between replicated points (since these edges do not affect our analysis, we ignore them for the sake of simplicity of exposition).

The graph G has average degree $O(M^5 \log^2 t) = O(\log^7 \mathcal{E})$, and the total weight of the vertices is $\Theta(\text{opt} \log \mathcal{E})$. Therefore, by Turán's theorem, one can compute an independent set of vertices in this graph of weight $\Omega(w(\mathbf{P}) / (\text{average degree} + 1)) = \Omega(\text{opt} / \log^6 \mathcal{E})$.

Finally, it is easy to verify that any triangle in \mathcal{T} contains at most 9 points of this independent set. ■

C.2 Covering a measure on a fat triangle

At this point we would like to use Lemma C.2 in order to get a bi-criteria approximation for the case in which the capacities are arbitrary, as we did in Theorem B.21. However, doing so directly proves more challenging for fat triangles than axis parallel rectangles. This is because our general procedure requires that, given an object x of a given type such that the total fractional value of the points in x is non-zero, we need to be able to decompose x into $O(\#(x))$ smaller objects of the same type such that, for each smaller object, the fractional value of the points in the object is only a constant. This can easily be done for axis parallel rectangles by using vertical splitting lines (as was done in Theorem B.21), but it is more challenging for fat triangles. However, the following lemma shows that such a decomposition is still possible for fat triangles.

Lemma C.3 *Let μ be a measure defined over the plane, and consider a fat triangle Δ . Then, for any integer k , one can cover Δ by at most $18k$ fat triangles, such that the measure of each of these triangles is at most $\mu(\Delta)/k$.*

Proof: To simplify the presentation we assume that $\mu(\Delta) = 1$. We recursively build a tree on Δ by partitioning the original triangle Δ into 4 similar triangles as shown in Figure 1. Each node of this tree corresponds to a triangle from this recursive construction. We stop the recursive partition for a node v as soon as the measure of the triangle Δ_v associated with it is at most $1/k$.

Once we have the tree, we select a set S of nodes of the tree as follows. We find the lowest node v in the tree such that the measure of its corresponding triangle is at least $1/k$. We add the node v to S and we treat the measure inside the triangle corresponding to v as being 0. We repeat this process until the measure left uncovered is smaller than $1/k$, at which point we take the lowest node covering the remaining measure and we add it to S . We also add the root of the tree to S . Note that the set S contains at most $k + 1$ nodes: since the initial measure is one, we added at most k nodes to S that are not the root.




Figure 1:

We also add all the nodes in this tree that are the least common ancestor (LCA) of a pair of nodes in S . Let S' be the resulting set of nodes. Now we can show that, for any tree T and any subset R of nodes of T , the set of all LCA nodes of all of the pairs of nodes in R has size at most $|R| - 1$ ¹¹. Therefore there are at most $|S| - 1$ nodes in $S' - S$ and thus the size of S' is at most $2|S| \leq 2(k + 1)$. Let \mathcal{T} be the set of triangles induced by the triangles corresponding to the children of the nodes of S' (i.e., every node of S' gives rise to four triangles). Consider the partition of the original triangle formed by \mathcal{T} . It is easy to verify that every face in this arrangement has measure at most $1/k$, and the number of faces of this arrangement, denoted by n , is at most $8k$ (observe that since the triangles arise out of a recursive partition, a pair of such triangles is either disjoint or contained in each other). Furthermore, each face of this arrangement is either a triangle (which is a scaled and rotated copy of the original triangle), or the difference of two triangles where one contains the other (having this property is why we took the children of every LCA). We will refer to a face which is the difference of two triangles as an annulus face. Let n' be the number of annulus faces in this arrangement. Clearly, $n' \leq |S'| = 2k$, as one can charge an annulus face to the node of S' that induced the hole in this face.




Figure 2:

We claim that an annulus face can be covered by the union of six translated and rotated copies of the original triangle. The easy case, is when the hole is a scaled and translated reflection of the outside triangle, see Figure 2, where three triangles are sufficient. The other case is when the hole is a scaled translated copy of the outer triangle, which by attaching to the hole three translated and rotated copies of the hole, gets reduced to the other case, see Figure 3. (The property used here implicitly is that the outer triangle of the annulus can be partitioned into translated and rotated copies of the hole triangle, as the hole arises out of a recursive partition of the outer triangle.)




Figure 3:

As such, there are $n - n'$ triangular faces in this arrangement and n' annulus faces. Thus, the original triangle can be covered by $(n - n') + 6n' = n + 5n' \leq 8k + 10k \leq 18k$ translated and scaled copies of the original triangle that cover it completely, and no triangle in this collection has measure that exceeds $1/k$. ■

Remark C.4 Given a weighted set of n points defining the measure inside the given fat triangle, the cover of Lemma C.3 can be computed in $O(n \log n)$ time. This requires using known techniques used in constructing compressed quadtrees, see [Har11] for details.

C.3 The result

Theorem C.5 *Given an instance of $(\mathcal{V}, \mathcal{T})$ of **PACKPNTSINFATTRIANGS** (with arbitrary capacities), one can compute, in polynomial time, a subset $X \subseteq \mathcal{V}$ that is $(O(\log^6 \mathcal{E}), 9)$ -approximation to the optimal solution.*

Proof: Compute a fractional solution to the given instance. For any triangle Δ in the plane, we denote by $\mathcal{E}(\Delta) = \sum_{p \in \mathcal{V} \cap \Delta} x_p$ the total mass of the fractional solution inside Δ . Next, we get a constant capacity instance out of $(\mathcal{V}, \mathcal{T})$ by replacing each triangle of \mathcal{T} by a “few” triangles covering

¹¹Let u, v be the pair of nodes in R whose LCA has maximum depth. Let z be the LCA of u and v , and let $R' = R - \{u, v\} \cup \{z\}$. The pairs in R and the pairs in R' have the same set of LCA nodes and thus the number of LCA nodes of a set of size r satisfies the recurrence $f(r) \leq f(r - 1) + 1$ and $f(2) = 1$.

it, such that the total mass of the fractional solution inside each of these new triangles is at most $c = 4 \cdot 18 \cdot 9$. Formally, consider a triangle $\Delta \in \mathcal{T}$, and let $k = \lceil \#(\Delta)/c \rceil$. If $\#(\Delta) \leq c$ then there is nothing to do (as $\mathcal{E}(\Delta) \leq \#(\Delta) \leq c$), so we assume that $\#(\Delta) > c$. Applying the algorithmic version of Lemma C.3, see Remark C.4, we cover Δ with at most

$$18k = 18 \left\lceil \frac{\#(\Delta)}{c} \right\rceil = 18 \left\lceil \frac{\#(\Delta)}{4 \cdot 18 \cdot 9} \right\rceil \leq \left\lceil \frac{\#(\Delta)}{2 \cdot 9} \right\rceil$$

triangles, where the total mass of the fractional solution inside each of them is at most $\mathcal{E}(\Delta)/k \leq \#(\Delta)/\lceil \#(\Delta)/c \rceil \leq c$.

Now, consider the generated instance with these new triangles, where each such triangle has capacity one. To this end, scale down the solution of the LP by a factor of c . Clearly, we now have a uniform capacity instance with an associated (valid) fractional solution having value $\Omega(\text{opt})$ (where opt is the optimal LP value for the original instance). Furthermore, any solution to this unit capacity instance, would correspond to a solution to the original instance (since we covered each original triangle with at most $18k \leq \left\lceil \frac{\#(\Delta)}{2 \cdot 9} \right\rceil \leq \#(\Delta)$ new unit capacity triangles). Plugging this instance into Lemma C.2 yields the required approximation. Specifically, every triangle $\Delta \in \mathcal{T}$ contains at most $9 \lceil \#(\Delta)/(2 \cdot 9) \rceil \leq \max(9, \#(\Delta))$ points of the computed set of points. ■

C.4 Canonical decomposition for fat triangles

In this section, we show that given a set \mathbf{P} of n points in the plane, and a parameter k , one can compute a set \mathcal{S} of $O(k^3 n \log^2 n)$ regions, such that for any α -fat triangle Δ , if $|\Delta \cap \mathbf{P}| \leq k$, then there exists (at most) 9 regions in \mathcal{S} whose union has the same intersection with \mathbf{P} as Δ does.

Our construction follows closely the argumentation of Aronov *et al.* [AES10]. However, our construction is (somewhat) different and (arguably) simpler since we are considering a “dual” problem to theirs. In particular, since modifying Aronov *et al.* [AES10] to get our result is not obvious, we present it here in detail.

C.4.1 Initial setup

To construct the set of regions, \mathcal{S} , we will use an approach similar to that of Lemma B.17. As was observed in [AES10], we can restrict our attention to axis aligned right triangles whose hypotenuse differs by no more than say one degree from -45° , as measured from the positive x -axis (i.e. it is near isosceles and faces to the right). In the following, let Δ be an arbitrary such triangle that contains at most k points.

We first construct a two level interval tree on P , where the first level partitions the points based on their x -coordinate, and the second level based on their y -coordinate (and the splitting line for each node goes through the median point). Let v be the highest node in the first level of the interval tree whose corresponding split line, ℓ , intersects Δ . Let Δ_{\leftarrow} and Δ_{\rightarrow} denote the portion of Δ to the left or right of ℓ , respectively. Also, let u be the highest node in the second level tree rooted at the left child of v whose corresponding split line, \wp , intersects Δ_{\leftarrow} , and let Δ_{\nwarrow} and Δ_{\swarrow} denote the portion of Δ_{\leftarrow} above or below \wp , respectively. (Note that we may assume that there exists split lines ℓ and \wp that intersect Δ and Δ_{\leftarrow} , respectively, since such regions that contain no points can be skipped). In the following, let p denote the point of intersection between ℓ and \wp . See Figure 4.




Figure 4: Decomposing Δ .

We now construct sets of canonical regions, $\mathcal{T}_{\rightarrow}$, \mathcal{T}_{\nwarrow} , and \mathcal{T}_{\swarrow} such that for any choice of Δ there exists constant number of regions r_1, \dots, r_m in $\mathcal{T}_{\rightarrow} \cup \mathcal{T}_{\nwarrow} \cup \mathcal{T}_{\swarrow}$, such that $\Delta \cap P = \bigcup_i (r_i \cap P)$, and $m \leq 9$.

We achieve this by showing that in each case (i.e., Δ_{\nwarrow} , Δ_{\swarrow} and Δ_{\rightarrow}) the region r under consideration can be transformed into a polygonal region with a constant number of points of P (or orientations) defining its bounding edges, and whose intersection with P is the same as r^{12} . We then show that the number of such regions needed for a particular choice ℓ and \wp is $O(nk^3)$.

In the following, let P_v be the subset of points of P stored in the subtree rooted v , and let $P_{u,v}$ be the set of points stored in the subtree rooted at u .

C.4.2 Handling the right portion of the triangle (Δ_{\rightarrow})

For any Δ , we know that Δ_{\rightarrow} will be a homothet of Δ , whose vertical edge lies on ℓ , see Figure 4. We now transform Δ_{\rightarrow} uniquely such that two points of P lie on its hypotenuse (or one point and the hypotenuse is at an angle of -46°) and one point of P lies on its bottom edge. Start by translating the hypotenuse towards the lower left corner of Δ_{\rightarrow} (while clipping it to Δ_{\rightarrow}) until it hits a point, p_1 . Next rotate the hypotenuse clockwise around p_1 until it hits a second point p_2 , or its orientation is -46° (as we rotate we modify its length so that one endpoint of the hypotenuse stays on ℓ and the other on the base of Δ_{\rightarrow}). Next translate the base of Δ_{\rightarrow} straight upwards (while clipping it and the hypotenuse as to maintain a right triangle) until it hits a third point p_3 (which may be the same as the rightmost point out of p_1 and p_2). Observe that the resulting region has the same intersection with P as Δ_{\rightarrow} (except maybe for the points on the boundary). See Figure 5.




Figure 5: Handling Δ_{\rightarrow} .

¹²For each point on the bounding edges, we will need to specify whether it is inside or outside the canonical region. This can be encoded by a string of length c , where c is some constant bounding the number of defining boundary points. Hence we can specify the inclusion or exclusion of the boundary points while only increasing the number of canonical regions by a factor of $2^c = O(1)$, and hence we will not need to worry about such issues.

Now we consider (what is by far) the hardest case, when $\mathbf{p} \notin \Delta_{\swarrow}$. In order to handle this case we will need to break up Δ_{\swarrow} as follows. Observe that Δ_{\swarrow} is a rectangular region whose upper right corner was cut off by the hypotenuse of Δ . First, we reduce Δ_{\swarrow} into a homothet of Δ , by removing rectangles r and r' from the left and bottom parts of Δ_{\swarrow} , respectively (see Figure 9). This can be done since we already observed that by the proof of Lemma B.17 we can construct a set of $O(nk^2 \log^2 n)$ canonical rectangles such that any rectangle (with a side on one of the split lines) has the same intersection with \mathbf{P} as one of the canonical rectangles. For simplicity we continue to refer to the remaining part of Δ_{\swarrow} as just Δ_{\swarrow} .




Figure 9:




Figure 10:

We now break up Δ_{\swarrow} into three regions. Let β^+ and β^- denote the rays emanating from \mathbf{p} at angles -140° and -130° , respectively (again, as measured clockwise from the positive x -axis). These two lines split Δ_{\swarrow} into three regions, which we will denote in their counterclockwise order as Δ_{\swarrow}^+ , Δ_{\swarrow}^0 and Δ_{\swarrow}^- (see Figure 10). Let a and b denote the intersection of β^+ with the hypotenuse and left edge of Δ_{\swarrow} , respectively. Similarly, let c and d denote the intersection of β^- with the hypotenuse and bottom edge of Δ_{\swarrow} , respectively.

Handling the top and bottom parts of Δ_{\swarrow} (i.e., Δ_{\swarrow}^+ and Δ_{\swarrow}^-). We now construct the canonical regions for Δ_{\swarrow}^+ . The construction is nearly identical to that for Δ_{\rightarrow} and is included for the sake of completeness. The construction for Δ_{\swarrow}^- is omitted as it is symmetric to the Δ_{\swarrow}^+ case.

Start by translating the part of the boundary that intersects the hypotenuse of Δ towards the lower left corner of Δ_{\swarrow}^+ (while clipping it to Δ_{\swarrow}^+) until it hits a point, \mathbf{p}_1 . Next rotate this edge counterclockwise around \mathbf{p}_1 until it hits a second point \mathbf{p}_2 , or its orientation is -44° (as we rotate we modify its length so that one endpoint stays on β^+ and the other on the boundary Δ_{\swarrow}^+). Next translate the vertical edge of Δ_{\swarrow}^+ to the right (while clipping it to Δ_{\swarrow}^+) until it hits a third point, \mathbf{p}_3 .




Figure 11:

As for the number of such resulting regions, assume that \mathbf{p}_2 lies to the left of \mathbf{p}_1 (the other case is handled similarly). There are $n_{u,v} = |\mathbf{P}_{u,v}|$ possible choices for \mathbf{p}_2 . Now consider the vertical line segment that connects β^+ and \mathbf{p}_2 . Imagine rotating this segment counterclockwise around \mathbf{p}_2 (while increasing its length so that the other endpoint stays on β^+) until it hits \mathbf{p}_1 . We know that all the points we hit in this sweeping process must be in our canonical region, and hence we can only have swept over k points before reaching \mathbf{p}_1 (if \mathbf{p}_2 does not exist we start with the triangle formed by β^+ and a vertical and -44° line through \mathbf{p}_1). Now imagine translating the vertical segment connecting \mathbf{p}_2 and β^+ to the left until we hit \mathbf{p}_3 (while increasing its length so that its top endpoint stays on the line defined by \mathbf{p}_1 and \mathbf{p}_2 and its bottom endpoint on β^+). Again we know that all the points we hit in this sweeping process must be in our canonical region, and hence we can only have swept over k points before reaching \mathbf{p}_3 . Hence there are $O(n_{u,v}k^2)$ such canonical regions for a pair nodes u and v . Thus overall there are $O(nk^2 \log^2 n)$ such canonical regions.

Handling the middle part of Δ_{\setminus} (i.e., Δ_{\setminus}^0).

Let $P_{u,v}^0$ denote the subset of $P_{u,v}$ that lies in between β^+ and β^- . Let the intersection of the hypotenuse of Δ with Δ_{\setminus}^0 be called the hypotenuse.

Translate the hypotenuse towards p (while clipping it to Δ_{\setminus}^0) until it hits a point p_1 . Then rotate the hypotenuse clockwise around p_1 until it hits a point p_2 , or it becomes vertical. Without loss of generality, assume that p_1 lies to the right of p_2 . Let the horizontal (resp. vertical) line connecting p_1 and β^+ (resp. β^-) be called h (resp. ν). Translate h downwards (resp. ν to the left), while enlarging it so that one endpoint stays on β^+ (resp. β^-), until either it hits the lowest (resp. furthest to the left) point of Δ_{\setminus}^0 or a point outside of Δ_{\setminus}^0 . Let this point be denoted p_3 (resp. p_4), and let h' (resp. ν') be the final translation of h (resp. ν). See Figure 12.




Figure 12:

Consider the region, r , bounded by the portion of h' to the left of p_4 , the portion of ν' below p_3 , β^+ , β^- , and the line going through p_1 and p_2 (this is the red shaded region in Figure 12). First observe that if both p_3 and p_4 lie outside of Δ_{\setminus}^0 then r will not cover all the points in $\Delta_{\setminus}^0 \cap P_{u,v}^0$. Namely, the points lying in the rectangle defined by h' , ν' , and the vertical and horizontal edges of Δ_{\setminus}^0 might not be covered by r (see Figure 12). However, we already constructed a set of $O(k^2 n \log n)$ canonical rectangles, which we know contains two canonical rectangles that cover these points, and as such we do not have to worry about covering these points. Clearly, all the points of $\Delta_{\setminus}^0 \cap P_{u,v}^0$ either lie in this rectangle or in r . Next observe that there are no points of $P_{u,v}^0$ that lie in r that are not in $\Delta_{\setminus} \cap P_{u,v}^0$. This follows from the easily proven fact (i.e. tedious but straightforward arguments) that since the hypotenuse was within one degree of -45° , that $h \cap \beta^+$ lies to the right of b and $\nu \cap \beta^-$ lies above d .

We now bound the number of canonical regions of type r . There are $|P_{u,v}^0|$ possible choices for p_1 (which again we assume is to the right of p_2). Now consider rotating h clockwise around p_1 until we hit p_2 . We know from above that all the points we sweep past in this process must be contained in $\Delta_{\setminus}^0 \cap P_{u,v}^0$ and so given p_1 there are at most k possible choices for p_2 . Now consider translating h downward (resp. ν to the left) until we hit p_3 (resp. p_4). Again, from above we know that all the points we sweep over in this process must be contained in $\Delta_{\setminus}^0 \cap P_{u,v}^0$ and so given p_1 and p_2 , there are at most k possible choices for p_3 (resp. p_4). Hence there are $O(k^3 |P_{u,v}^0|)$ such canonical regions for a given pair of nodes u and v , and so overall there are $O(k^3 n \log^2 n)$ such canonical regions.

C.4.5 Putting things together

Summing the above bounds over all choices of the nodes u and v results overall in $O(k^3 n \log^2 n)$ canonical regions. Furthermore, for any choice of Δ , we showed above that there exists a set of at most 9 of these canonical regions whose (union of) intersections with P is the same as that of Δ . We thus get the following result.

Theorem C.6 *Given a set P of n points in the plane, and parameters k and $\alpha > 0$, one can*

compute a set \mathcal{S} of $O(k^3 n \log^2 n)$ regions, such that for any α -fat triangle Δ , if $|\Delta \cap \mathcal{P}| \leq k$, then there exists (at most) 9 regions in \mathcal{S} whose union has the same intersection with \mathcal{P} as Δ does.

D PTAS for Unweighted Disks and Points

In this section, we consider instances of the **PACKREGIONS** problem in which the regions are disks with unit weights and all points have unit capacities. We now outline a PTAS for such instances based on the local search technique. The algorithm and proof are an extension of those of Chan and Har-Peled [CH11, CH09], and Mustafa and Ray [MR10].

The algorithm. Since all of the regions have unit weight, we may assume that no region is completely contained in another. We say that a subset L of \mathcal{D} is ***b -locally optimal*** if L is a pointwise independent set and one cannot obtain a larger pointwise independent set by removing $\ell \leq b$ regions of L and inserting $\ell + 1$ regions of $\mathcal{D} \setminus L$.

Our algorithm constructs a b -locally optimal solution using local search, where b is some suitable constant. We start with $L \leftarrow \emptyset$. We consider each subset $X \subseteq \mathcal{D} \setminus L$ of size at most $b + 1$: if X is a pointwise independent set and the set $Y \subseteq L$ of regions pointwise intersecting the objects of X has size at most $|X| - 1$, we set $L \leftarrow (L \setminus Y) \cup X$. Every such swap increases the size of L by at least one, and as such it can happen at most $n = |\mathcal{D}|$ times. Therefore the running time is bounded by $O(n^{b+3} b |\mathcal{P}|)$, since there are $\binom{n}{b+1}$ subsets X to consider and for each such subset X it takes $O(nb |\mathcal{P}|)$ time to compute Y .

Analysis. Let opt be the maximum pointwise independent set, and let L be the b -locally optimal solution returned by our algorithm. If we can show that the pointwise intersection graph of $\text{opt} \cup L$ is planar then the analysis in [CH11] will directly imply that $|L| \geq (1 - O(1/\sqrt{b})) |\text{opt}|$.

We map the disks in opt and L to sets of points Q_{opt} and Q_L in \mathbb{R}^3 , respectively, and we map the points in \mathcal{P} to a set of halfspaces $H_{\mathcal{P}}$, by using the lifting of disks to planes and points to rays, and then dualizing the problem (see Appendix B.2). Mustafa and Ray prove that a range space defined by a set of points and halfspaces in \mathbb{R}^3 has the ***locality condition***, which is defined as follows.

Definition D.1 ([MR10]) *A range space $R = (\mathcal{P}, \mathcal{D})$ satisfies the **locality condition** if for any two disjoint subsets $R, B \subseteq \mathcal{P}$, it is possible to construct a planar bipartite graph $G = (R, B, E)$ with all edges going between R and B such that for any $D \in \mathcal{D}$, if $D \cap R \neq \emptyset$ and $D \cap B \neq \emptyset$, then there exist two vertices $u \in D \cap R$ and $v \in D \cap B$ such that $(u, v) \in E$.*

Since opt and L are both pointwise independent sets, we know each point in \mathcal{P} can intersect at most one disk from opt and at most one disk from L . Hence each halfspace in $H_{\mathcal{P}}$ can contain at most one point from Q_{opt} and at most one point from Q_L . Since points and halfspaces in \mathbb{R}^3 have the locality condition, setting $R = L$ and $B = \text{opt}$ immediately implies that there is a planar graph on the vertex set $L \cup \text{opt}$ such that any vertex from L and any vertex from opt that are in the same halfspace are adjacent. In particular, the intersection graph is planar.

Theorem D.2 *Given a set of n unweighted disks and a set of m points in the plane (with unit capacities), any b -locally optimal pointwise independent set has size $\geq (1 - O(1/\sqrt{b})) \text{opt}$, where opt is the size of the maximum pointwise independent set of the disks. In particular, one can compute an independent set of size $\geq (1 - \varepsilon) \text{opt}$, in time $mn^{O(1/\varepsilon^2)}$.*

Corollary D.3 *There is a PTAS for instances of **PACKHALFSPACES** in which each halfspace has unit weight, and each point has unit capacity.*

Corollary D.4 *There is a PTAS for instances of **PACKREGIONS** in which each region is a unit-weight disk, and each point has unit capacity.*

Corollary D.5 *There is a PTAS for instances of **PACKPOINTS** in which each region is a unit-capacity disk, and each point has unit weight.*

E Hardness of approximation


E.1 Packing same size fat triangles into points

Here we show that **PACKREGIONS** (Problem 1.2) does not have a PTAS, even if the regions have unit weight and their union complexity is linear. We show that the problem is APX-hard using a reduction from the maximum bounded 3-dimensional matching problem. Since maximum bounded 3-dimensional matching is APX-complete [Kan91], this will imply the claim (unless $P = NP$).

Theorem E.1 *Unless $P = NP$ there is no PTAS for **PACKREGIONS** (Problem 1.2) even if the regions are unweighted, in the plane, and have linear union complexity. In particular, this holds if the regions are fat triangles of similar size. (See Corollary B.3 (A) for the matching approximation algorithm.)*

Proof: Let $T \subseteq A \times B \times C$ be the input triples for an instance of maximum bounded 3-dimensional matching, where A , B , and C are disjoint subsets of some ground set X (for simplicity we assume $X = A \cup B \cup C$). For each element $x \in X$ we make a representative point v_x and place it arbitrarily on the unit circle in the plane and give it unit capacity. Let V_A , V_B , and V_C be the sets of representatives for A , B , and C (respectively). A triple in T thus corresponds to a triangle with one vertex in each of V_A , V_B , and V_C . Clearly, finding a maximum packing of these triangles into these points is an instance of **PACKREGIONS**. Moreover, a maximum packing here corresponds to a maximum set of triangles (triples) such that each point (element of X) is covered by at most one triangle. Therefore a PTAS for this problem translates to a PTAS for the maximum bounded 3-dimensional matching problem. (Note that this does not imply that there is no PTAS for other specific types of regions.)

Now we show that we can make the triangles fat and of similar size, and hence there is no PTAS even in the case of linear union complexity. Let the range of a set of representative points be the angle around the circle between the farthest two points of the set, and let the center of a set be the midpoint on the circle between the farthest two points of the set. Instead of placing the points arbitrarily, we will place the points so that the range of each of V_A , V_B , or V_C is less than five degrees. Moreover, we place the points so that the centers of V_A , V_B , and V_C are 120 degrees apart. In this case the triangles all have roughly the same size and are nearly equilateral. It is known that such a set of triangles has linear union complexity [MMP⁺94]. Hence, by the above reduction, even in this case where the regions are restricted to have linear union complexity (and even more specifically when they are restricted to be fat triangles of roughly the same size), we cannot get a PTAS. ■



E.2 Packing points into fat triangles

Lemma E.2 *There is an approximation-preserving reduction from the **INDEPENDENT SET** problem in general graphs to the **PACKPOINTS** problem. In particular, for instances of the problem **PACKPOINTS** in which the regions are fat triangles with unit capacities and the points are unweighted, no approximation better than $\Omega(n^{1-\varepsilon})$ is possible in polynomial time, for any constant $\varepsilon > 0$, unless $P = NP$.*

Proof: Consider an instance of the **INDEPENDENT SET** problem, namely a graph $G = (V, E)$. Let $n = |V|$. Place n distinct points on the unit circle (arbitrarily) and map every vertex of V to a unique point of the resulting set of points P . For every edge $uv \in E$, consider the segment $p_u p_v$, where p_u and p_v are the points corresponding to u and v in P . We construct a fat triangle containing $p_u p_v$ by connecting p_u , p_v , and a third vertex in the interior of the unit disk; this can always be done so as to achieve roughly 2-fatness. We add this triangle to our set of regions \mathcal{D} , and assign it capacity one.

Clearly, solving the resulting instance (P, \mathcal{D}) of **PACKPOINTS** is equivalent to solving the **INDEPENDENT SET** problem for G . The claim now follows from the hardness results known for the **INDEPENDENT SET** problem [Has99]. \blacksquare

F Additional details

F.1 Basic tools

The following lemma testifies that the packing problem can be solved in a straightforward fashion if all the capacities are the same (i.e., uniform capacities). This is done by repeatedly applying a procedure to find and remove an independent set in the remaining induced sub-hypergraph (for example the procedure in [CH11]).

Lemma F.1 *Let G be a hypergraph for which there is a polynomial time algorithm `alg` that takes as input a fractional solution to **HYPERGRAPH-LP** for an **HGRAPHPACKING** instance on G , or any induced subgraph of G , with unit capacities — i.e., an independent set instance — and it constructs an integral solution whose value is at least an α fraction of the value of the fractional solution. Then one can compute in polynomial time a 2α -approximation for any instance of **HGRAPHPACKING** on G with uniform capacities (i.e., all the hyperedges have the same capacity, say k).*

Proof: Let $G = (V, E)$ be an instance of **HGRAPHPACKING** in which all hyperedges have the same capacity, say k . Let $G_0 = G$, and in the i th iteration, for $i = 1, \dots, k$, compute a maximum weight independent set Y_i in G_{i-1} using `alg`, and let $G_i = G_{X \setminus U_i}$, where $U_i = Y_1 \cup \dots \cup Y_i$. We claim that U_k is the required approximation.

Clearly, no hyperedge of G contains more than k vertices of U_k as it is the union of k independent sets, and as such it is a valid solution. Now, let V_{opt} be the optimal solution. If $w(V_{\text{opt}} \cap U_k) \geq w(V_{\text{opt}})/2$ then we are done. Otherwise, consider the hypergraph G_{i-1} , and observe that $V_{\text{opt}} \setminus U_{i-1}$ is a valid solution for **HGRAPHPACKING** for this graph (with uniform capacities k). Interpreting this integral solution as a solution to the LP, and scaling it down by k , we get a fractional solution to the independent set LP of this hypergraph of value $w(V_{\text{opt}} \setminus U_{i-1})/k$. Since Y_i was constructed using `alg` on the optimal fractional solution to the independent set LP of this hypergraph, we have that

$$w(Y_i) \geq \text{opt}_{\text{LP}}(G_{i-1})/\alpha \geq w(V_{\text{opt}} \setminus U_{i-1})/k\alpha \geq w(V_{\text{opt}} \setminus U_k)/k\alpha \geq w(V_{\text{opt}})/2k\alpha.$$

Which implies that $w(U_k) \geq w(V_{\text{opt}})/2\alpha$. \blacksquare

A hypergraph $G = (V, E)$ *shatters* $X \subseteq V$ if the number of hyperedges in G_X is $2^{|X|}$. The **VC dimension** of G is the size of the largest set of vertices it shatters.

The following is a “sparsification” lemma. Here we get better bounds than the standard technique, as we are using stronger sampling results known for spaces with bounded VC dimension.

Lemma F.2 *Let $G = (V, E)$ be an instance of **HGRAPHPACKING** with VC dimension d , and consider its fractional LP solution of value opt and with energy \mathcal{E} . Then, one can compute, in polynomial time, a valid fractional solution for the LP of G such that:*

- (A) *The value of the new fractional solution is $\geq \text{opt}/12$.*
- (B) *The number of vertices with non-zero value is $O(d\mathcal{E} \log \mathcal{E})$.*
- (C) *The value of each non-zero variable is equal to i/M for some integer $i \leq M$, where $M = O(d \log \mathcal{E})$.*
- (D) *The total energy in the new solution is $\Theta(\mathcal{E})$.*

Proof: Let $\varepsilon = 1/\mathcal{E}$, where $\mathcal{E} = \sum_v x_v$, and x_v is the value the LP assigns to $v \in V$ in the optimal LP solution. Let $T = O(d \log \mathcal{E})$ and R be a random sample of V of (expected) size $\tau = \mathcal{E}T = O((d/\varepsilon) \log(1/\varepsilon))$, created by picking each vertex v independently with probability $x_v \cdot T$ ¹³. This sample is a relative $(\varepsilon, 1/2)$ -approximation [Har11, HS11], with probability of failure $\leq \rho_1 = \varepsilon^{O(d)}$. That implies that for any hyperedge $f \in E$ such that $x(f) = \sum_{v \in f} x_v$ we have

$$\frac{|R \cap f|}{|R|} \leq (1 + 1/2) \left(\frac{x(f)}{\mathcal{E}} + \varepsilon \right).$$

To interpret the above, observe that $\mathbf{E}[|R \cap f|] = x(f)T$ and $\mathbf{E}[|R|] = \mathcal{E}T$, as such, a rough estimate of the expectation of $|R \cap f| / |R|$ is $x(f)/\mathcal{E}$. Thus, the above states (somewhat opaquely) that no hyperedge is being over-sampled by R .

Since the expected size of R is τ , by Chernoff’s inequality, we know that $|R| \leq 2\tau$ with probability at least $1 - \rho_2$, where $\rho_2 = \varepsilon^{O(d\mathcal{E})}$ (as $\mathcal{E} \geq 1$). Now, consider a hyperedge f with capacity k , and observe that $x(f) \leq k$. As such, $|R \cap f| \leq (1 + 1/2)(x(f)/\mathcal{E} + \varepsilon) |R| \leq (3/2)(k + 1)\varepsilon 2\tau \leq 6kT$. In particular, if v appears t_v times in R (R is a multiset), then we assign it the fractional value $y_v = t_v/6T$. We then have that $y(f) = \sum_{v \in f} y_v \leq |R \cap f|/6T \leq k$ (and this holds for all hyperedges with probability $\geq 1 - \rho_1$). As such, the fractional solution defined by the y ’s is valid.

As for the value of this fractional solution, consider the random variable $Z = \sum_v y_v w(v)$, which is a function of the random sample R . Observe that

$$\mathbf{E}[Z] = \sum_v w(v) \mathbf{E}\left[\frac{t_v}{6T}\right] = \sum_v \frac{w(v)}{6T} \mathbf{E}[t_v] = \sum_v \frac{w(v)}{6T} x_v T = \frac{1}{6} \sum_v w(v) x_v = \frac{\text{opt}_{\text{LP}}}{6},$$

as $\text{opt}_{\text{LP}} = \sum_v w(v) x_v$. In particular, since no vertex can have $w(v) > \text{opt}_{\text{LP}}$ (otherwise, we would choose it as the solution), it follows that

$$Z = \sum_v y_v w(v) \leq \text{opt}_{\text{LP}} \sum_v \frac{t_v}{6T} \leq \text{opt}_{\text{LP}} \frac{|R|}{6T} \leq \text{opt}_{\text{LP}} \frac{2\tau}{6T} = \text{opt}_{\text{LP}} \frac{2\mathcal{E}T}{6T} \leq \frac{\mathcal{E}}{3} \text{opt}_{\text{LP}}.$$

¹³A minor technicality is that $x_v T$ might be larger than one. In this case, we put $\lfloor x_v T \rfloor$ copies of v into R , and we put an extra copy of v into R with probability $x_v T - \lfloor x_v T \rfloor$. It is straightforward to verify that our argumentation goes through in this case. Observe that such large values work in our favor by decreasing the probability of failure.

This implies that $\Pr\left[Z \geq \text{opt}_{\text{LP}}/12\right] \geq 1/4\varepsilon = \varepsilon/4$. Indeed, if not,

$$\begin{aligned} \mathbf{E}\left[Z\right] &\leq \frac{\text{opt}_{\text{LP}}}{12} \Pr\left[Z \leq \frac{\text{opt}_{\text{LP}}}{12}\right] + \Pr\left[Z \geq \frac{\text{opt}_{\text{LP}}}{12}\right] \frac{\varepsilon}{3} \text{opt}_{\text{LP}} \\ &< \frac{\text{opt}_{\text{LP}}}{12} + \frac{1}{4\varepsilon} \cdot \frac{\varepsilon}{3} \text{opt}_{\text{LP}} \leq \frac{\text{opt}_{\text{LP}}}{6}, \end{aligned}$$

a contradiction. As such, a random sample R corresponds to a valid assignment with value at least $\text{opt}_{\text{LP}}/12$ with probability at least $\Pr[Z \geq \text{opt}_{\text{LP}}/12] - \rho_1 - \rho_2 \geq \varepsilon/8$, as $\rho_1 + \rho_2$ is an upper bound on the sample R failing to have the desired properties. As such, taking $u = O(\varepsilon \log \varepsilon)$ independent random samples one of them is the required assignment, with probability $\geq 1 - (1 - \varepsilon/8)^u \geq 1 - 1/\varepsilon^{O(1)}$. We take this good sample together with its associated LP values as the desired fractional solution to the LP. Also, note that the total energy of the new solution is $\Theta(\varepsilon)$, since by Chernoff's inequality $\tau/2 \leq |R| \leq 2\tau$ with probability at least $1 - 2\rho_2$. \blacksquare

F.2 Improving the running time

In Section 3, we described an algorithm that constructs an ordering of the vertices by repeatedly finding the vertex of least resistance with respect to the set of remaining vertices. Computing the resistance of a vertex by brute force takes $O(n^{C+O(1)})$ time, where C is the maximum capacity of an edge in E . However, for our analysis to go through, we only need to find a vertex that is safe with respect to the set of remaining vertices; informally, a vertex v is safe if the probability that it participates in a conflict with a random sample of the remaining vertices is smaller than some constant (that is strictly smaller than one), where each remaining vertex u is included in the sample with probability x_u/Δ . In this section we show that there is a sampling algorithm that finds a safe vertex with high probability and its running time is polynomial in the maximum capacity C .

Lemma F.3 *Computing a good ordering of the vertices can be done in polynomial time. Namely, the algorithm of Section 3 can be implemented in polynomial time.*

Proof: To get the same quality of approximation we do not need to take the vertex of least resistance in each round (of computing the ordering), but merely a vertex that is “safe.” More precisely, let X be the current set of vertices, let v be a vertex of this set, and let R be a random sample of X in which each vertex u is included with probability x_u/Δ (also we force v to be in R). We say that v is **violated** in R if v is contained in a hyperedge f such that the number of vertices of f that are in R is larger than its capacity $\#(f)$. Let $\mu(v, X)$ denote the probability that v is violated in R . Note that $\mu(v, X)$ is a (conservative) upper bound on the probability that v is rejected by our rounding algorithm if we started with an ordering in which $X \setminus \{v\}$ is the set of all vertices that come before v . Therefore, in order for our rounding to succeed, in each round we only need to find a vertex v for which the probability $\mu(v, X)$ is low, where X is the set of all vertices that still need to be ordered at the beginning of the round. (We remark that it follows from the argument of Lemma 3.9 that, for any set X , there is a vertex v for which $\mu(v, X) \leq 1/4$.)

Now we are ready to describe how to construct an ordering for our algorithm. Let X be the set of vertices that still need to be ordered. As we will see shortly, for each vertex $v \in X$, we can compute an estimate $\bar{\mu}(v, X)$ of the probability $\mu(v, X)$. We pick the vertex v with minimum estimated probability $\bar{\mu}(v, X)$, we make v the last vertex (in the ordering of X) and we recursively order $X \setminus \{v\}$.

We can compute the estimates $\bar{\mu}(v, X)$ in polynomial time as follows. Fix a vertex v . Let ψ be a sufficiently large polynomial in n . We pick ψ independent random samples of X (again, forcing v

to be in each of these samples); in each random sample, each vertex u is included with probability x_u/Δ . We set $\bar{\mu}(v, X)$ to be the fraction of the samples in which the vertex v is violated. Using a standard argument based on the Chernoff inequality, we can show that our estimates are very close with high probability, and therefore our rounding algorithm achieves the required approximation with high probability as well; we omit the easy but tedious details. ■

G Proofs

G.1 Proof of Lemma 3.2

Proof: Let $X_i = \langle v_1, \dots, v_i \rangle$. Note that, if we selected v_i , we rejected v_i in the alteration phase only if v_i participates in a conflict with some of the vertices in $\{v_1, \dots, v_{i-1}\} \cap \mathcal{C}$. Let Z_i be the number of conflicts of X_i that contain v_i and are realized in \mathcal{C} , i.e., $h \subseteq \mathcal{C}$. In the following, we show that the probability that Z_i is non-zero is at most $1/4$, which implies the lemma.

Consider a k -conflict $h = \{v_{j_1}, \dots, v_{j_k}, v_i\}$, where each vertex of h is in X_i and h contains v_i . The probability that all of the vertices of h are selected, given that v_i is selected, is equal to $\frac{x_{j_1}}{\Delta} \cdot \frac{x_{j_2}}{\Delta} \dots \frac{x_{j_k}}{\Delta} = \frac{\Delta}{x_i} \rho_\Delta(h)$. Therefore we have

$$\mathbf{E}\left[Z_i \mid v_i \in \mathcal{C}\right] = \sum_{h \in \mathcal{H}, h \subseteq X_i, v_i \in h} \frac{\Delta}{x_i} \rho_\Delta(h) = \eta_\Delta(v_i, X_i) \leq \frac{1}{4},$$

where the last inequality follows from Lemma 3.9 and the fact that v_i is the vertex of minimum resistance in X_i . Thus

$$\Pr\left[v_i \notin \mathcal{O} \mid v_i \in \mathcal{C}\right] \leq \Pr\left[Z_i > 0 \mid v_i \in \mathcal{C}\right] \leq \mathbf{E}\left[Z_i \mid v_i \in \mathcal{C}\right] \leq \frac{1}{4}.$$

Therefore, if v_i is selected, it is accepted with probability at least $3/4$. ■

G.2 Proof of Corollary 3.3

Proof: By Lemma 3.2, for each vertex $v \in \mathcal{V}$, we have

$$\begin{aligned} \Pr\left[v \in \mathcal{O}\right] &= \Pr\left[(v \in \mathcal{O}) \cap (v \in \mathcal{C})\right] = \Pr\left[v \in \mathcal{O} \mid v \in \mathcal{C}\right] \Pr\left[v \in \mathcal{C}\right] \geq \frac{3}{4} \Pr\left[v \in \mathcal{C}\right] \\ &\geq \frac{3x_v}{4\Delta}, \end{aligned}$$

where $\Delta = O(\gamma(\mathcal{E})^{1/\nu})$. By linearity of expectation, we have that the expected weight of the generated solution is at least

$$\sum_{v \in \mathcal{V}} \frac{3x_v}{4\Delta} w_v = \Omega\left(\frac{\sum_v x_v w_v}{\gamma(\mathcal{E})^{1/\nu}}\right) = \Omega\left(\frac{\text{opt}}{\gamma(\mathcal{E})^{1/\nu}}\right),$$

as $\sum_v x_v w_v$ is the value of the fractional LP solution, which is bigger than (or equal to) the weight of the optimal solution. ■

G.3 Proof of Lemma 3.4

Proof: Each k -conflict h that is realized corresponds to a hyperedge f with capacity k such that $h = f \cap \mathbf{R}$. Additionally, two realized conflicts that are distinct correspond to different hyperedges. Therefore the number of k -conflicts that are realized in \mathbf{R} is at most the number of hyperedges f such that the capacity of f is k and $|f \cap \mathbf{R}| = k + 1$. It follows from the definition of $F_k(\cdot)$ that the number of k -conflicts is at most $F_k(|\mathbf{R}|)$. Therefore it suffices to upper bound the expected value of $F_k(|\mathbf{R}|)$.

Note that $\mathbf{E}[|\mathbf{R}|] = \mathcal{E}(X)/2$. We have

$$\begin{aligned} \mathbf{E}\left[F_k(|\mathbf{R}|)\right] &\leq \sum_{t=0}^{\infty} \Pr\left[|\mathbf{R}| \geq t \frac{\mathcal{E}(X)}{2}\right] F_k\left(\left(t+1\right) \frac{\mathcal{E}(X)}{2}\right) \leq \sum_{t=0}^{\infty} 2^{-(t+1)/2} F_k\left(\left(t+1\right) \frac{\mathcal{E}(X)}{2}\right) \\ &\leq \sum_{t=0}^{\infty} 2^{-(t+1)/2} c^{O(\log t)} F_k\left(\mathcal{E}(X)\right) = O\left(F_k\left(\mathcal{E}(X)\right)\right), \end{aligned}$$

since \mathbf{G} has the bounded growth property (see Definition 2.1), and by the Chernoff inequality (we use here implicitly that $\mathcal{E}(X) \geq 1$). ■

G.4 Proof of Lemma 3.5

Proof: Let $f \in \mathbf{E}$ be a hyperedge with capacity k that generated the conflict h . Since x is a feasible solution for the LP, we have that $\sum_{v \in f-h} x_v \leq \sum_{v \in f} x_v \leq \#(f) = k$. Clearly, the conflict h is realized if we pick all the vertices of h , and none of the vertices of $f - h$, and the probability of that event is

$$\begin{aligned} \prod_{v \in h} \frac{x_v}{2} \prod_{v \in f-h} \left(1 - \frac{x_v}{2}\right) &\geq \frac{1}{2^{k+1}} \prod_{v \in h} x_v \prod_{v \in f-h} \exp(-x_v) \\ &= \frac{\rho(h)}{2^{k+1}} \cdot \exp\left(-\sum_{v \in f-h} x_v\right) \geq \frac{\rho(h)}{2(2e)^k}, \end{aligned}$$

In the first line we used the inequality $1 - x_v/2 \geq \exp(-x_v)$, which holds since $x_v \leq 1$. ■