Math 2415

Problem Section #10

15.1 Double Integrals over Rectangles

- 1. Evaluate the double integral, $\iint_R (2y+3) dA$, where $R = [0,3] \times [0,2]$ by identifying it as the volume of a solid. (No need to compute an integral!)
- 2. Calculate $\iint_R y e^{xy} dA$ where $R = [0, 1] \times [0, 2]$.
- 3. (a) Explain why the paraboloid $z = 2 + x^2 + y^2$ always lies under the plane z = 8 when (x, y) is in the rectangle $[0, 1] \times [0, 2]$.
 - (b) Sketch the solid enclosed by the paraboloid $z = 2+x^2+y^2$ and the planes x = 0, x = 1, y = 0, y = 2, and z = 8. **Hint:** Do this by sketching the curves obtained by intersecting $z = 2 + x^2 + y^2$ in the four planes x = 0, x = 1, y = 0, and y = 2.
 - (c) Set up a double integral to calculate the volume of this solid and evaluate the integral.

15.2: Double Integrals (Rectangular Coordinates)

- 1. Sketch a region that is Type I but not Type II.
- 2. Set up iterated integrals for both orders of integration for the integral $\iint_D y \, dA$, where *D* is bounded by x = 0, y = x and y = 3 x. In which order is easier to do the iterated integrals? Explain. Evaluate the integral this way.
- 3. Evaluate $\iint_D x^2 dA$ where D is the triangular region with vertices (0, 2), (1, 3), and (4, 0).
- 4. Evaluate the integral, $\int_{x=0}^{x=1} \int_{y=x^2}^{y=1} \sqrt{y} \sin(y) \, dy \, dx$ by reversing the order of integration.
- 5. Find the volume of the tetrahedron bounded by the coordinate planes and the plane x + 2y + 3z = 6.
- 6. (a) Explain why the plane z = x always lies under the the plane z = 4 over the region in the *xy*-plane between $y = x^2$ and $y = 1 x^2$.
 - (b) Find the volume of the solid region under the plane z = 4, above the plane z = x, and between the parabolic cylinders $y = x^2$ and $y = 1 x^2$.

Exam Two Review

You can start review for Exam Two today. We will do more of these next week.

- 1. Spring 2019: 3
- 2. Fall 2017: 6
- 3. Fall 2016 Exam II: 1,2,4,6,7
- 4. Fall 2014 Exam II: 1,2,3,4
- 5. Fall 2012 Exam II:1,2,3,4,6,8