

# Math 2415

## Problem Section #7

**Make sure you do some problems from each section.**

### 14.3-14.5: Partial Derivatives, Tangent Planes, and Chain Rule

**Chain Rule:** Let  $z = f(x, y)$  be a function on the plane and let  $(x, y) = \mathbf{r}(t)$  be a curve in the plane. The composition

$$z = g(t) = f(\mathbf{r}(t)) = f(x(t), y(t))$$

is a scalar-valued function of one variable. The function  $g$  is called the **restriction** of  $f$  to the curve  $\mathbf{r}$ , since it just gives us the values of  $f$  along the curve  $\mathbf{r}$ . In this context, the **Chain Rule for Functions on Curves** states that

$$g'(t) = \nabla f(\mathbf{r}(t)) \cdot \mathbf{r}'(t) = f_x(x(t), y(t))x'(t) + f_y(x(t), y(t))y'(t).$$

*Now for the questions!*

- Let  $f(x, y) = xy + x^2 = x(y + x)$  Calculate  $f_x$ ,  $f_y$ ,  $f_{xy}$  and  $f_{xx}$  at the point  $\mathbf{x}_0 = (1, 1)$ . Discuss the geometric meaning of each of these derivatives, with the aid of schematic diagrams showing slices of  $f$  in planes where  $x$  or  $y$  are constant.
- Show that the function  $u(x, y) = e^{-x} \cos(y)$  satisfies Laplace's equation  $u_{xx} + u_{yy} = 0$ .
- Show that the function  $u(x, t) = \cos(kx) \sin(akt)$  satisfies the wave equation  $u_{tt} = a^2 u_{xx}$ .
- Find an equation of the form  $z = Ax + By + C$  for the tangent plane to the function  $z = f(x, y) = e^x \cos(xy)$  at  $(x_0, y_0) = (0, 0)$ . Explain why your solution shows that  $e^x \cos(xy) \approx x + 1$  near  $(0, 0)$ .
- Let  $z = f(x, y) = y^2 \sin x$  where  $(x, y) = \mathbf{r}(t) = (e^{3t}, t^4)$ .
  - Form the composition  $g(t) = f(x(t), y(t))$  and then use the single variable chain rule to calculate  $g'(t)$ .
  - Use the Chain Rule for Functions on Curves to calculate  $g'(t)$ .
- Use the chain rule to calculate  $\frac{\partial z}{\partial s}$  and  $\frac{\partial z}{\partial t}$  where  $z = \cos(x^2 + y^2)$  and  $x = t \ln s$ ,  $y = se^t$ . Hint: Make a tree diagram showing the relationships between the variables.
- Let  $g(t) = f(x(t), y(t))$ , where  $x(2) = 6$ ,  $x'(2) = 8$ ,  $y(2) = -1$ ,  $y'(2) = 3$ ,  $f(6, -1) = 10$ ,  $f_x(6, -1) = 2$ ,  $f_y(6, -1) = 7$ ,  $f(8, 3) = -4$ ,  $f_x(8, 3) = 5$ ,  $f_y(8, 3) = 9$ . Find  $g'(2)$ .
- Suppose that  $z = f(x, y)$  and that  $g(u, v) = f(\cos(u) + v^2, \sin(u) - v^3)$ . Use the table of values to calculate  $g_u(0, 1)$  and  $g_v(0, 1)$ .

$(x, y)$	$f$	$f_x$	$f_y$
$(0, 1)$	5	3	-7
$(2, -1)$	3	9	-4

- The temperature at point  $(x, y)$  on a hot plate is  $T = T(x, y)$ . An ant walks on the hot plate so that its position at time  $t$  is  $x = 1 + t^2$ ,  $y = t^3$ . If  $\nabla T(5, 8) = (6, -1)$  find the rate of change of the ant's temperature at time  $t = 2$ .

## 16.6, Parametrized Surfaces

1. Let  $S$  be the surface with parametrization

$$(x, y, z) = \mathbf{r}(u, v) = u \cos v \mathbf{i} + u \sin v \mathbf{j} + u \mathbf{k} \quad u \geq 0, \quad 0 \leq v \leq 2\pi.$$

- (a) Show that  $S$  is a cone. **Hint:** Find an equation of the form  $F(x, y, z) = 0$  for this surface by eliminating  $u$  and  $v$  from the equations for  $x$ ,  $y$ , and  $z$  above.
- (b) Sketch the cone, together with the “grid” curves on the cone where (a)  $u = 2$  and (b)  $v = \pi/4$ .
- (c) Find a parametrization of the tangent plane to the cone at the point where  $(u, v) = (2, \pi/4)$ . Add this tangent plane to your sketch.
2. (a) Write down the equation of the form  $F(x, y, z) = 0$  for the sphere of radius 2, center  $(1, 2, 3)$ .
- (b) Show that

$$(x, y, z) = \mathbf{r}(\theta, \phi) = (1 + 2 \sin \phi \cos \theta, 2 + 2 \sin \phi \sin \theta, 3 + 2 \cos \phi)$$

is a parametrization of this sphere. **Hint:** Substitute the formulae for  $x$ ,  $y$ , and  $z$  in terms of  $\theta$  and  $\phi$  into the function  $F$  you obtained in (a) and simplify as much as you can. What does this calculation tell you about where each of the points  $(x, y, z) = \mathbf{r}(\theta, \phi)$  lie?

3. Find a parametrization for each of the following surfaces. [Note: There are many correct answers!] Show how you arrived at your answer. **Hint:** It is often helpful to construct your parametrization using (a) cylindrical coordinates, (b) spherical coordinates, or (c) by using a parametrization such as  $(x, y, z) = (u, v, f(u, v))$  for the surface  $z = f(x, y)$ .
- (a) The portion of the paraboloid  $z = x^2 + y^2$  where  $z \leq 4$ .
- (b) The portion of the cone  $z = 2\sqrt{x^2 + y^2}$  that is between the planes  $z = 2$  and  $z = 4$  and is in the first octant.
- (c) The portion of the sphere  $x^2 + y^2 + z^2 = 9$  that is above the cone  $z = \sqrt{x^2 + y^2}$ .
- (d) The portion of the cylinder  $y^2 + z^2 = 9$  between the planes  $x = 0$  and  $x = 3$ .