



Statistical Methods in AI and ML

Nicholas Ruozzi

University of Texas at Dallas

A powerful and flexible set of tools for modeling
problems in AI/ML

Judea Pearl won the Turing award for his work on
Bayesian networks!
(among other achievements)

Exploit **locality** and structural features of a given model in order to gain insight about **global properties**

- What this course is:
 - Probabilistic graphical models
 - Topics:
 - representing data
 - exact and approximate statistical inference
 - model learning
 - variational methods in ML

- What you should be able to do at the end:
 - Design statistical models for applications in your domain of interest
 - Apply learning and inference algorithms to solve real problems (exactly or approximately)
 - Understand the complexity issues involved in the modeling decisions and algorithmic choices


- CS 5343: Algorithm Analysis and Data Structures
- CS 3341: Probability and Statistics in Computer Science and Software Engineering
- Basically, comfort with probability and algorithms (machine learning is helpful, but not required)

Suggested Textbook



Readings will be posted
online before each
lecture

Check the course website
for additional resources
and papers



- In addition, some lecture notes, in book format, will be made available for the main topics
- The idea is to build a set of notes that aligns well with the presentation of course material
- Comments, suggestions, corrections are welcome/encouraged

- 4-6 problem sets (80%)
 - See collaboration policy on the web
- Lecture scribes (15%)
- Class/Piazza participation & extra credit (5%)

-subject to change-

Course Info.



- Instructor: Nicholas Ruozzi
 - Office: ECSS 3.409
 - Office hours: TBD, and by appointment
- TA: TBD
 - Office hours and location TBD
- Course website:
<http://www.utdallas.edu/~nrr150130/cs6347/2024sp/>

- Model the world (or at least the problem) as a collection of random variables related through some joint probability distribution
 - Compactly represent the distribution
 - Undirected graphical models
 - Directed graphical models
- Learn the distribution from observed data
 - Maximum likelihood, SVMs, etc.
- Make predictions (statistical inference)

Inference and Learning




```
107 51 108 51 109 31 110 41 111 51 112 41 113 51 11 41 114 51 115 51 116 51 1
132 51 133 51 134 51 13 51 135 31 136 41 137 51 138 51 139 41 140 51 141 41 1
31 156 21 157 51 158 41 159 51 160 21 161 51 162 51 163 21 16 41 164 51 165 4
181 51 182 11 183 41 184 51 18 51 185 51 186 11 187 41 188 31 189 41 190 31 1
21 206 51 207 51 208 31 209 51 210 41 211 41 212 31 21 31 213 41 214 51 215 4
231 41 232 11 23 31 233 51 234 21 24 11 2 51 25 51 26 51 27 11 28 51 29 21 30
31 48 51 49 21 50 21 51 51 52 51 53 31 54 11 5 51 55 31 56 51 57 51 58 51 59
31 77 51 78 31 79 41 80 51 81 41 82 51 8 31 83 51 84 41 85 21 86 51 87 21 88
52 241 32 242 52 243 42 244 52 245 52 246 52 247 52 248 42 249 52 250 52 251
32 268 22 269 52 270 32 271 52 272 52 273 52 274 52 275 52 276 42 277 52 278
52 295 42 296 33 297 53 298 53 299 43 300 33 301 53 302 33 303 53 304 53 305
54 322 44 323 54 324 44 325 54 326 44 327 44 328 54 329 54 330 44 331 24 332
44 349 54 350 54 351 54 352 54 353 24 354 44 355 55 356 55 357 55 358 55 359
45 376 45 377 55 378 35 379 45 380 55 381 45 382 45 383 55 384 55 385 55 386
56 403 57 213 47 232 47 386 47 387 57 404 57 405 57 406 47 407 57 408 47 409
37 426 57 427 57 428 57 429 47 430 37 431 57 432 47 433 47 434 47 435 47 436
47 453 47 454 57 455 47 456 37 457 47 458 47 459 47 460 47 461 47 462 47 463
47 480 47 481 47 482 47 483 57 484 47 485 57 486 57 487 57 488 47 489 47 490
47 507 57 508 37 509 47 510 47 511 47 512 47 513 47 514 47 515 47 516 47 517
57 534 57 535 27 536 47 537 47 538 47 539 47 540 47 541 57 542 37 543 47 544
37 561 47 562 47 563 47 564 57 565 47 566 47 567 47 568 47 569 47 570 47 571
47 588 57 589 47 590 47 591 47 592 27 593 47 594 47 595 47 596 37 597 47 598
47 615 47 616 47 617 47 618 47 619 47 620 48 621 38 622 48 623 18 624 58 625
48 642 48 643 48 644 48 645 38 646 38 647 49 296 49 528 59 648 49 649 59 650
49 667 59 668 59 669 49 670 49 671 59 672 49 673 59 674 19 675 49 676 59 677
49 694 59 695 49 696 49 697 29 698 59 699 59 700 59 701 29 702 59 703 59 704
49 721 59 722 59 723 59 724 29 725 59 726 59 727 59 728 59 729 59 730 49 731
```

Collect Data

$$Z(\theta) = \sum_x p(x; \theta)$$

Use the model to do inference / make predictions



“Learn” a model that represents the observed data

Inference and Learning




```
107 51 108 51 109 31 110 41 111 51 112 41 113 51 11 41 114 51 115 51 116 51 1
132 51 133 51 134 51 13 51 135 31 136 41 137 51 138 51 139 41 140 51 141 41 1
31 156 21 157 51 158 41 159 51 160 21 161 51 162 51 163 21 16 41 164 51 165 4
181 51 182 11 183 41 184 51 18 51 185 51 186 11 187 41 188 31 189 41 190 31 1
21 206 51 207 51 208 31 209 51 210 41 211 41 212 31 21 31 213 41 214 51 215 4
231 41 232 11 23 31 233 51 234 21 24 11 2 51 25 51 26 51 27 11 28 51 29 21 30
31 48 51 49 21 50 21 51 51 52 51 53 31 54 11 5 51 55 31 56 51 57 51 58 51 59
31 77 51 78 31 79 41 80 51 81 41 82 51 8 31 83 51 84 41 85 21 86 51 87 21 88
52 241 32 242 52 243 42 244 52 245 52 246 52 247 52 248 42 249 52 250 52 251
32 268 22 269 52 270 32 271 52 272 52 273 52 274 52 275 52 276 42 277 52 278
52 295 42 296 33 297 53 298 53 299 43 300 33 301 53 302 33 303 53 304 53 305
54 322 44 323 54 324 44 325 54 326 44 327 44 328 54 329 54 330 44 331 24 332
44 349 54 350 54 351 54 352 54 353 24 354 44 355 55 356 55 357 55 358 55 359
45 376 45 377 55 378 35 379 45 380 55 381 45 382 45 383 55 384 55 385 55 386
56 403 57 213 47 232 47 386 47 387 57 404 57 405 57 406 47 407 57 408 47 409
37 426 57 427 57 428 57 429 47 430 37 431 57 432 47 433 47 434 47 435 47 436
47 453 47 454 57 455 47 456 37 457 47 458 47 459 47 460 47 461 47 462 47 463
47 480 47 481 47 482 47 483 57 484 47 485 57 486 57 487 57 488 47 489 47 490
47 507 57 508 37 509 47 510 47 511 47 512 47 513 47 514 47 515 47 516 47 517
57 534 57 535 27 536 47 537 47 538 47 539 47 540 47 541 57 542 37 543 47 544
37 561 47 562 47 563 47 564 57 565 47 566 47 567 47 568 47 569 47 570 47 571
47 588 57 589 47 590 47 591 47 592 27 593 47 594 47 595 47 596 37 597 47 598
47 615 47 616 47 617 47 618 47 619 47 620 48 621 38 622 48 623 18 624 58 625
48 642 48 643 48 644 48 645 38 646 38 647 49 296 49 528 59 648 49 649 59 650
49 667 59 668 59 669 49 670 49 671 59 672 49 673 59 674 19 675 49 676 59 677
49 694 59 695 49 696 49 697 29 698 59 699 59 700 59 701 29 702 59 703 59 704
49 721 59 722 59 723 59 724 29 725 59 726 59 727 59 728 59 729 59 730 49 731
```

Data sets can be large

$$Z(\theta) = \sum_x p(x; \theta)$$

Inference needs to be fast



Data must be compactly modeled

Applications



- Computer vision
- Natural language processing
- Robotics
- Computational biology
- Computational neuroscience
- Text translation
- Text-to-speech
- Many more...

- A graphical model is a graph together with "local interactions"
- The graph and interactions model a global optimization or learning problem
- The study of graphical models is concerned with how to exploit local structure to solve these problems either exactly or approximately



Probability Review

- **Sample space** specifies the set of possible outcomes
 - For example, $\Omega = \{H, T\}$ would be the set of possible outcomes of a coin flip
- Each element $\omega \in \Omega$ is associated with a number $p(\omega) \in [0,1]$ called a **probability**

$$\sum_{\omega \in \Omega} p(\omega) = 1$$

- For example, a biased coin might have $p(H) = .6$ and $p(T) = .4$

- An **event** is a subset of the sample space
 - Let $\Omega = \{1, 2, 3, 4, 5, 6\}$ be the 6 possible outcomes of a dice roll
 - $A = \{1, 5, 6\} \subseteq \Omega$ would be the event that the dice roll comes up as a one, five, or six
- The probability of an event is just the sum of all of the outcomes that it contains
 - $p(A) = p(1) + p(5) + p(6)$

Independence




- Two events A and B are **independent** if

$$p(A \cap B) = p(A)p(B)$$

Let's suppose that we have a fair die: $p(1) = \dots = p(6) = 1/6$

If $A = \{1, 2, 5\}$ and $B = \{3, 4, 6\}$ are A and B independent?



Independence




- Two events A and B are **independent** if

$$p(A \cap B) = p(A)p(B)$$

Let's suppose that we have a fair die: $p(1) = \dots = p(6) = 1/6$

If $A = \{1, 2, 5\}$ and $B = \{3, 4, 6\}$ are A and B independent?




No!

$$p(A \cap B) = 0 \neq \frac{1}{4}$$

Independence




- Now, suppose that $\Omega = \{(1,1), (1,2), \dots, (6,6)\}$ is the set of all possible rolls of two **unbiased** dice
- Let $A = \{(1,1), (1,2), (1,3), \dots, (1,6)\}$ be the event that the first die is a one and let $B = \{(1,6), (2,6), \dots, (6,6)\}$ be the event that the second die is a six
- Are A and B independent?



Independence



- Now, suppose that $\Omega = \{(1,1), (1,2), \dots, (6,6)\}$ is the set of all possible rolls of two **unbiased** dice
- Let $A = \{(1,1), (1,2), (1,3), \dots, (1,6)\}$ be the event that the first die is a one and let $B = \{(1,6), (2,6), \dots, (6,6)\}$ be the event that the second die is a six
- Are A and B independent?



- The **conditional probability** of an event A given an event B with $p(B) > 0$ is defined to be

$$p(A|B) = \frac{p(A \cap B)}{P(B)}$$

- This is the probability of the event $A \cap B$ over the sample space $\Omega' = B$
- Some properties:
 - $\sum_{\omega \in B} p(\omega|B) = 1$
 - If A and B are independent, then $p(A|B) = p(A)$

Simple Example



Cheated	Grade	Probability
Yes	A	.15
Yes	F	.05
No	A	.5
No	F	.3

Simple Example



Cheated	Grade	Probability
Yes	A	.15
Yes	F	.05
No	A	.5
No	F	.3

$$p(\text{Cheated} = \text{Yes} \mid \text{Grade} = \text{F}) = ?$$


Simple Example




Cheated	Grade	Probability
Yes	A	.15
Yes	F	.05
No	A	.5
No	F	.3

$$p(\textit{Cheated} = \textit{Yes} \mid \textit{Grade} = \textit{F}) = \frac{.05}{.35} \approx .14$$


The Monty Hall Problem



1



2



3

$$p(A \cap B) = p(A)p(B|A)$$

$$\begin{aligned} p(A \cap B \cap C) &= p(A \cap B)p(C|A \cap B) \\ &= p(A)p(B|A)p(C|A \cap B) \end{aligned}$$

·
·
·

$$p\left(\bigcap_{i=1}^n A_i\right) = p(A_1)p(A_2|A_1) \dots p(A_n|A_1 \cap \dots \cap A_{n-1})$$

Conditional Independence



- Two events A and B are independent if learning something about B tells you nothing about A (and vice versa)
- Two events A and B are **conditionally independent** given C if

$$p(A \cap B|C) = p(A|C)p(B|C)$$

- This is equivalent to

$$p(A|B \cap C) = p(A|C)$$

- That is, given C , information about B tells you nothing about A (and vice versa)

Conditional Independence



- Let $\Omega = \{(H, H), (H, T), (T, H), (T, T)\}$ be the outcomes resulting from tossing two different fair coins
- Let A be the event that the first coin is heads
- Let B be the event that the second coin is heads
- Let C be the event that both coins are heads or both are tails
- A and B are independent, but A and B are not independent given C

- A discrete **random variable**, X , is a function from the state space Ω into a discrete space D

- For each $x \in D$,

$$p(X = x) \equiv p(\{\omega \in \Omega : X(\omega) = x\})$$

is the probability that X takes the **value** x

- $p(X)$ defines a probability distribution
 - $\sum_{x \in D} p(X = x) = 1$
- Random variables partition the state space into disjoint events

Example: Pair of Dice



- Let Ω be the set of all possible outcomes of rolling a pair of dice
- Let p be the uniform probability distribution over all possible outcomes in Ω
- Let $X(\omega)$ be equal to the sum of the value showing on the pair of dice in the outcome ω
 - $p(X = 2) = ?$
 - $p(X = 8) = ?$

Example: Pair of Dice



- Let Ω be the set of all possible outcomes of rolling a pair of dice
- Let p be the uniform probability distribution over all possible outcomes in Ω
- Let $X(\omega)$ be equal to the sum of the value showing on the pair of dice in the outcome ω
 - $p(X = 2) = \frac{1}{36}$
 - $p(X = 8) = ?$

Example: Pair of Dice



- Let Ω be the set of all possible outcomes of rolling a pair of dice
- Let p be the uniform probability distribution over all possible outcomes in Ω
- Let $X(\omega)$ be equal to the sum of the value showing on the pair of dice in the outcome ω
 - $p(X = 2) = \frac{1}{36}$
 - $p(X = 8) = \frac{5}{36}$

- We can have vectors of random variables as well

$$X(\omega) = [X_1(\omega), \dots, X_n(\omega)]$$

- The **joint distribution** is $p(X_1 = x_1, \dots, X_n = x_n)$ is

$$p(X_1 = x_1 \cap \dots \cap X_n = x_n)$$

typically written as

$$p(x_1, \dots, x_n)$$

- Because $X_i = x_i$ is an event, all of the same rules - independence, conditioning, chain rule, etc. - still apply

- Two random variables X_1 and X_2 are independent if

$$p(X_1 = x_1, X_2 = x_2) = p(X_1 = x_1)p(X_2 = x_2)$$

for all values of x_1 and x_2

- Similar definition for conditional independence
- The conditional distribution of X_1 given $X_2 = x_2$ is

$$p(X_1 | X_2 = x_2) = \frac{p(X_1, X_2 = x_2)}{p(X_2 = x_2)}$$

this means that this relationship holds for all choices of x_1

Expected Value



- The **expected value** of a real-valued random variable is the weighted sum of its outcomes

$$E[X] = \sum_{x \in D} p(X = d) \cdot d$$

- Expected value is linear

$$E[X + Y] = E[X] + E[Y]$$

Expected Value: Lotteries



- Powerball Lottery currently has a grand prize of \$102 million
- Odds of winning the grand prize are $1/292,201,338$
- Tickets cost \$2 each
- Expected value of the game

$$= \left(\frac{292,201,337}{292,201,338} \right) (-2) + \left(\frac{1}{292,201,338} \right) \cdot (102,000,000 - 2)$$

$$\approx \$ - 1.65$$

- The **variance** of a random variable measures its squared deviation from its mean

$$\mathit{var}(X) = E[(X - E[X])^2]$$

- Estimates the square of the expected amount by which a random variable deviates from its expected value