

4. Let $f(x) = x^{-3}(x - \sin x)$ for $x \neq 0$. How should $f(0)$ be defined in order that f be continuous? Will it also be differentiable?
5. a. Derive the Taylor series at 0 for the function $f(x) = \ln(x + 1)$. Write this series in summation notation. Give two expressions for the remainder when the series is truncated.
- b. Determine the smallest number of terms that must be taken in the series to yield $\ln 1.5$ with an error less than 10^{-8} .
- c. Determine the number of terms necessary to compute $\ln 1.6$ with error 10^{-10} at most.
6. Determine whether the following function is continuous, once differentiable, or twice differentiable:

$$f(x) = \begin{cases} x^3 + x - 1 & \text{if } x \leq 0 \\ x^3 - x - 1 & \text{if } x > 0 \end{cases}$$

7. (Continuation) Repeat the preceding problem for the function

$$f(x) = \begin{cases} x & \text{if } x \leq 1 \\ x^2 & \text{if } x > 1 \end{cases}$$

8. Criticize this reasoning: The function f defined by

$$f(x) = \begin{cases} x^3 + x & \text{if } x \leq 0 \\ x^3 - x & \text{if } x \geq 0 \end{cases}$$

has the properties

$$\lim_{x \rightarrow 0^+} f''(x) = \lim_{x \rightarrow 0^+} 6x = 0$$

$$\lim_{x \rightarrow 0^-} f''(x) = \lim_{x \rightarrow 0^-} 6x = 0$$

Therefore, f'' is continuous.

9. Prove that if f is differentiable at x , then

$$\lim_{h \rightarrow 0} \frac{f(x+h) - f(x-h)}{2h} = f'(x)$$

Show that for some functions that are not differentiable at x , the preceding limit exists. (See Eggermont [1988] or the following problem.)

10. Prove or disprove this assertion: If f is differentiable at x , then for $\alpha \neq 1$,

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x+\alpha h)}{h - \alpha h}$$

11. Show that $\lim_{x \rightarrow 1} (4x + 2) = 6$ by means of an ε - δ proof.
12. Show that $\lim_{x \rightarrow 2} (1/x) = \frac{1}{2}$ by means of an ε - δ proof.
13. For the function $f(x) = 3 - 2x + x^2$ and the interval $[a, b] = [1, 3]$, find the number ξ that occurs in the Mean-Value Theorem.
14. (Continuation) Repeat the preceding problem with the function $f(x) = x^6 + x^4 - 1$ and the interval $[0, 1]$.
15. Find the Taylor series for $f(x) = \cosh x$ about the point $c = 0$.
16. If the series for $\ln x$ is truncated after the term involving $(x - 1)^{1000}$ and is then used to compute $\ln 2$, what bound on the error can be given?

17. Find the Taylor series for $f(x) = e^x$ about the point $c = 3$. Then simplify the series and show how it could have been obtained directly from the series for f about $c = 0$.
18. Let k be a positive integer and let $0 < \alpha < 1$. To what classes $C^n(\mathbb{R})$ does the function $x^{k+\alpha}$ belong?
19. Prove: If $f \in C^n(\mathbb{R})$, then $f' \in C^{n-1}(\mathbb{R})$ and $\int_a^x f(t) dt$ belong to $C^{n+1}(\mathbb{R})$.
20. Prove Rolle's Theorem directly (not as a special case of the Mean-Value Theorem).
21. Prove: If $f \in C^n(\mathbb{R})$ and $f(x_0) = f(x_1) = \dots = f(x_n) = 0$ for $x_0 < x_1 < \dots < x_n$, then $f^{(n)}(\xi) = 0$ for some $\xi \in (x_0, x_n)$. *Hint:* Use Rolle's Theorem n times.
22. Prove that the function $f(x) = x^2$ is continuous everywhere.
23. For small values of x , the approximation $\sin x \approx x$ is often used. Estimate the error in using this formula with the aid of Taylor's Theorem. For what range of values of x will this approximation give results correct to six decimal places?
24. For small values of x , how good is the approximation $\cos x \approx 1 - \frac{1}{2}x^2$? For what range of values will this approximation give correct results rounded to three decimal places?
25. Use Taylor's Theorem with $n = 2$ to prove that the inequality $1 + x < e^x$ is valid for all real numbers except $x = 0$.
26. Derive the Taylor series with remainder term for $\ln(1+x)$ about 1. Derive an inequality that gives the number of terms that must be taken to yield $\ln 4$ with error less than 2^{-m} .
27. What is the third term in the Taylor expansion of $x^2 + x - 2$ about the point 3?
28. Using the series for e^x , how many terms are needed to compute e^2 correctly to four decimal places (rounded)?
29. Develop the Taylor series for $f(x) = \ln x$ about e , writing the results in summation notation and giving the remainder term. Suppose $|x - e| < 1$ and accuracy $\frac{1}{2} \times 10^{-1}$ is desired. What is the minimum number of terms in the series required to achieve this accuracy?
30. Determine the first two terms of the Taylor series for x^x about 1 and the remainder term E_1 .
31. Determine the Taylor polynomial of degree 2 for $f(x) = e^{(\cos x)}$ expanded about the point π .
32. First develop the function \sqrt{x} in a series of powers of $(x - 1)$ and then use it to approximate $\sqrt{0.9999999995}$ to ten decimal places.
33. Assume that $|x| < \frac{1}{2}$ and determine by Taylor's Theorem the best upper bound.
- $|\cos x - (1 - x^2/2)|$
 - $|\sin x - x(1 - x^2/6)|$
34. Determine a function that can be termed the **linearization** of $x^3 - 2x$ at 2.
35. How many terms are required in the series

$$e = \sum_{k=0}^{\infty} \frac{1}{k!}$$

to give e with an error of at most $6/10$ unit in the 20th decimal place?

36. Find the first two terms in the Taylor expansion of $x^{1/5}$ about the point $x = 32$. Approximate the fifth root of 31.999999 using these two terms in the series. How accurate is your answer?
37. Find the Taylor polynomial of degree 2 for the function $f(x) = e^{2x} \sin x$ expanded about the point $\pi/2$.